

Mixed Dentition Treatment with Cervical Traction and Lower Lingual Arch

WILLIAM M. ODOM

Dr. Odom is Associate Clinical Professor of Orthodontics at the University of the Pacific in San Francisco, California, and in the private practice of Orthodontics in San Mateo and Half Moon Bay. He holds a D.D.S. degree from the College of Physicians and Surgeons in San Francisco, and a Certificate in Orthodontics from the University of Oregon.

A cephalometric comparison of treatment changes among cervical traction alone, cervical traction combined with banded upper incisors and a lower lingual arch, and untreated controls.

LITERATURE REVIEW OF MIXED DENTITION TREATMENT OF CLASS II MALOCCLUSION

There is little question that desirable treatment results can be achieved with a wide variety of appliances when they are used by skilled clinicians. Nevertheless, no clearly superior force system has come to the fore, and little commonality of opinion exists regarding treatment of Class II malocclusions at any stage of development.

The suggestion that treatment ought to begin in the mixed dentition is at least as old as Angle.¹⁵ A considerable body of literature has been published extolling the advantages of early treatment.¹⁻¹⁷ Mathews¹⁵ has suggested that his colleagues "preferentially" undertake the treatment of Class II malocclusions in the late deciduous or very early mixed dentition stage.

Maxillary Dentition

Many previous mixed dentition studies have evaluated the effects of extraoral traction in the treatment of

Address:

Dr. William Odom
University of the Pacific
Department of Orthodontics
2155 Webster Street
San Francisco, California

Class II malocclusions in the mixed dentition.

Following the reintroduction of extraoral traction by Kloehn,¹⁸ the orthodontic literature blossomed with numerous and often conflicting reports on its effects on the dentition, maxilla, mandible and associated structures.¹⁷⁻⁴³

Wieslander,¹⁷ studying the effect of cervical traction on both early and late mixed dentition samples, found treatment results in the early mixed dentition to be more favorable.

King,¹⁶ after reflecting on the success and failure of over 20 years of mixed dentition treatment, ends his paper expressing the hope that more study and research will expand the possibilities for definitive early treatment of Class II malocclusions.

It should be expected that the diversity of approaches used to treat such a diverse population as the seductively simple classification of Class II malocclusion would produce superficially conflicting reports.

Extraoral traction has been reported to produce varying effects on the occlusal plane,^{4,20-28,37} a frequent tendency for downward tipping of the palatal plane,^{20,22-24,27} an extrusive effect on the upper first molar,^{25-28,34,37,40} an increase in vertical face height^{28-31,34} and mandibular plane angle,^{34-36,40} and reduced forward movement of pogonion.^{21,24,27,28,34}

On the other hand, none of those effects occur universally with any of the many different force systems in use.³⁵⁻³⁸

Taking each of the diverse revelations of the different addenda to the literature as gospel would invariably lead to the condemnation or acceptance of some approach or other, even though such a posture might not be generally supported by the greater

body of literature. In this light, the recently published works of Baumrind *et al*⁴⁰⁻⁴² provide a new perspective by which we can reevaluate some old beliefs.

Various investigators have consistently reported a retarding effect on the forward growth of the maxilla^{13,22,24,27,43} or distal movement of upper teeth,^{20,22,26,28} sometimes with a change in the position of the pterygo-maxillary fissure^{22,35} and the sphenoid bone.^{24,35,36} Considering the combined findings of animal and clinical studies⁴² on the effects of applying a distally directed force to the maxilla, one is led to the inevitable conclusion that the forward growth of the maxilla can be retarded and that an absolute distal displacement of the maxilla and maxillary dentition is possible.

Mandibular Arch

The lingual archwire has been used successfully in many ways over the years. A concise discussion has been published by Singer.⁴⁴ This versatile appliance has numerous applications, but the apparent simplicity of its design belies the complexity of its mastery. As expressed by Singer, "Certainly, it can be seen that the appellation 'passive lingual arch' is a misnomer."

MATERIALS AND METHODS

It is easy to criticize previous investigations on the basis of their subjectivity, sample size, lack of a control sample, case report method, failure to control variables, statistics and so on.³⁴ Viewed from a broader perspective, each of these flawed attempts has added information which contributes to our better understanding in a speciality that is still heavily dependent on the art with which mechanical systems are applied.

TABLE 1

Samples

	Number			Age (Yr.) Film #1		Age (Yr.) Film #2		Elapsed Time (Yr.)
	Total	M	F	Mean	S.D.	Mean	S.D.	
Control	23	11	12	8.75	1.19	12.72	1.29	3.97
Treatment	21	12	9	8.54	2.1	12.51	2.3	3.97
KC Only	19	10	9	9.88	1.65	12.97	1.64	3.09

The observations of Moyers and Bookstein⁴⁵ notwithstanding, the data presented in this study was obtained by a retrospective analysis of lateral cephalometric radiographs of paired subjects who had been clinically diagnosed in the early mixed dentition as having a Class II malocclusion.

The treatment sample (Table 1) consisted of 21 individuals, 12 male and 9 female, each of whom received treatment for correction of the antero-posterior dysplasia, establishment of normal overbite and overjet relationships, maintenance of mandibular leeway space and alignment of the lower anterior teeth. It was not expected that a second phase of treatment would be required in the permanent dentition.

The control sample consisted of 11 males and 12 females who also presented with similar Class II malocclusions in the early mixed dentition.

An additional treatment sample of 10 males and 9 females, referred to as "KC Only," received treatment with only a Kloehe cervical traction appliance for the correction of the antero-posterior dysplasia.

The experimental sample was selected solely on the basis of pre-treatment records and criteria. Sample selection was approached in a manner as close as possible to statistical randomness, without conscious bias. No

consideration was given to the treatment result.

The University of California Combined Head Film Analysis⁴⁶ was applied to all radiographs. This method, developed by Dr. Sheldon Baumrind, is a computer-aided procedure which refines the precision with which films can be compared.

The 21 dental and skeletal landmarks shown in Fig. 1 were located on each film. Paired films were compared with registrations on the palatal plane, anterior cranial base and mandibular border.

Each film was traced by four different individuals, producing four independent estimates for each landmark and anatomic superimposition. Each estimate of each point was digitized on a two-dimensional coordinate system and averaged to yield a best estimate for the coordinates of each point. Additional computer operations superimposed those average patterns on each of the three anatomic reference bases.

Treatment

All subjects in the experimental sample were treated with a partial Edgewise appliance augmented by cervical traction and a lower lingual arch. The maxillary incisors and all four first molars were banded.

Treatment in the maxillary arch




Fig. 1 Landmarks used in U.C.S.F. Combined Headfilm Analysis.

- | | | |
|-----------------------|------------------------|---------------------------|
| 1. Porion | 8. Lower Incisor Edge | 15. Lower Molar Cusp |
| 2. Sella | 9. Point B | 16. Lower Molar Apex |
| 3. Nasion | 10. Lower Incisor Apex | 17. Upper Gonion Image |
| 4. Orbitale | 11. Pogonion | 18. Lower Gonion Image |
| 5. Upper Incisor Apex | 12. Menton | 19. Condyle |
| 6. Point A | 13. Upper Molar Apex | 20. Anterior Nasal Spine |
| 7. Upper Incisor Edge | 14. Upper Molar Cusp | 21. Posterior Nasal Spine |

These data were acquired using hardware and software technologies developed at the Craniofacial Research Instrumentation Laboratory of the Department of Growth and Development at the University of California, San Francisco, under NIH-NIDR Grants DE03598 and DE03703.

was initiated with a multistranded twisted .0215" (0.55mm) archwire, followed by .020" (0.51mm) round and .019" $\times$.025" (0.48 $\times$ 0.64mm) rectangular archwires with vertical offsets distal to the lateral incisors. Compression coil springs were used between the first molars and lateral incisors to maintain or gain space for the eruption of the permanent cuspids and bicuspids.

In some cases deciduous teeth were extracted to gain more favorable eruption of the permanent cuspids.

Cervical traction was applied to the first molars with a .045" (1.14mm) Kloehe type facebow. The long outer bows were raised to prevent distal crown tipping of the molars and possible incisor extrusion.

In the mandibular arch a symmetrical preformed .030" (0.77mm) lingual arch with a vertical loop mesial to the horizontal molar sheath was used. The lingual arch was adjusted to contact the lower incisors only, near the level

of their contact points. The arch form was modified as required for cuspid eruption.

The objectives of this phase of treatment were a normal molar relationship, normal overbite and overjet, well-aligned and positioned incisors and adequate space for the eruption of permanent cuspids and bicuspids.

When those goals were achieved, the maxillary incisor bands were removed and a removable palatal appliance with an anterior elastic was worn full-time. During this supervision/retention phase the cervical appliance was worn as required to maintain molar relationships.

The second records were made to evaluate the need for further treatment after the second molars had begun to erupt. Of the 21 individuals in the treatment sample, 16 were judged to not require additional banded treatment.

The results are shown in Figs. 2-11 and Tables 2-21.




Fig. 2 Upper first molar (U6) eruption during treatment averaged 0.25 mm/yr more than the controls.

TABLE 2
Treatment vs. Controls
Vertical Changes in Position of U6 (mm)

	Control		Treatment		dX	t
	Mean	S.D.	Mean	S.D.		
to Anterior Cranial Base	7.3	1.3	8.4	3.2	1.1	1.50
to S-N	7.15	1.4	8.5	2.8	1.35	2.05
to Palatal Plane	2.15	1.0	3.1	1.7	0.95	2.26*

* Significant at 0.05 level of confidence.




Fig. 3 Experimental treatment had a significant mean orthopedic effect, inhibiting the forward displacement of the upper first molar with growth with no significant change in dento-alveolar relationships.

TABLE 3
Treatment vs. Controls
Horizontal Changes in Position of U6

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
to Anterior Cranial Base	3.9	1.5	0.7	3.8	4.6	4.71**
to S-N	4.20	1.85	0.0	3.7	4.2	4.77**
to Palatal Plane	2.05	1.5	2.4	2.3	0.35	0.57

** Significant at 0.01 level of confidence.




Fig. 4 The mean effects of KloeHN headgear alone on the first molar were primarily distal movement within the alveolar process.

TABLE 4
Treatment vs. Cervical
Horizontal Changes in Position of U6

	<i>Treatment</i>		<i>KC</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
to Anterior Cranial Base	0.7	4.6	0.0	2.6	0.7	0.63
to S-N	0.0	4.2	0.3	2.2	0.3	0.66
to Palatal Plane	2.4	2.3	0.7	1.5	1.70	2.93**

** Significant at 0.01 level of confidence.




Fig. 5 Experimental treatment had no significant effect on restricting the mean downward growth of the upper central incisors.

TABLE 5
Treatment vs. Control
Vertical Changes in Position of U1

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
to Anterior Cranial Base	6.45	1.75	7.2	3.0	0.75	0.69
to S-N	6.35	1.80	7.2	2.7	0.85	1.81
to Palatal Plane	1.55	1.05	1.7	1.4	0.15	1.67




Fig. 6 Experimental treatment had a significant mean orthopedic effect in inhibiting the forward growth of the maxillary anterior alveolar process and central incisors.

TABLE 6
Treatment vs. Control
Horizontal Changes in Position of U1

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
	<i>Mesial</i>		<i>Distal</i>			
to Anterior Cranial Base	2.4	2.7	—3.2	4.7	3.6	4.87**
to S-N	2.75	2.7	—2.9	4.4	5.65	5.14**
to Palatal Plane	0.45	2.2	—0.7	2.4	2.85	2.82*

* Significant at 0.05 level of confidence.

** Significant at 0.01 level of confidence.




Fig. 7 Differences in the mean horizontal effects of experimental treatment and Kloehe cervical traction on the horizontal position of the upper central incisors were not statistically significant.

TABLE 7
Treatment vs. Cervical
Horizontal Changes in Position of U1

	<i>Treatment</i>		<i>KC</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
	<i>Distal</i>		<i>Distal</i>			
to Anterior Cranial Base	-3.2	4.7	-2.0	2.9	1.2	1.17
to S-N	-2.9	4.4	-1.6	2.6	1.3	1.57
to Palatal Plane	-.07	2.4	-1.4	2.2	0.7	0.875




Fig. 8 Experimental treatment, including lower lingual arch, did not inhibit lower molar eruption.

TABLE 8
Treatment vs. Control
Vertical (Upward) Changes in Position of L6

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
to Anterior Cranial Base	6.7	1.9	8.8	3.1	2.1	2.73*
to Mandibular Plane	2.65	1.35	4.1	1.8	1.45	3.02**

* Significant at 0.05 level of confidence.

** Significant at 0.01 level of confidence.




Fig. 10 Lower lingual arch did not prevent forward growth of the lower first molar, but did result in a more upright angulation than in the controls.

TABLE 9
Treatment vs. Cervical
Vertical (Upward) Changes in Position of L6

	<i>Treatment</i>		<i>KC</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
to Anterior Cranial Base	8.8	3.1	6.9	3.0	1.9	1.86
to Mandibular Plane	4.1	1.8	2.6	1.9	1.5	2.68*

* Significant at 0.05 level of confidence.



Fig. 9 More mean lower molar eruption occurred with experimental treatment including lower lingual arch than with Kloehe cervical therapy alone.

TABLE 10
Treatment vs. Control
Horizontal Changes in Position L6

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
to Anterior Cranial Base	3.9	1.5	-0.2	3.8	4.71	4.71**
Crown to Mandibular Plane	1.5	1.55	1.7	1.7	0.2	0.41
Apex to Mandibular Plane	0.4	1.55	3.0	3.3	2.6	3.38**
Tip (degrees)	1.85	4.45	-4.6	8.6	6.45	3.15

** Significant at 0.01 level of confidence.




Fig. 11 The lower li
gual arch did not signi
cantly affect the erupti
or angulation of the low
incisors.

TABLE 11
Treatment vs. Control
Changes in Position of L1

	Control		Treatment		dX	t
	Mean	S.D.	Mean	S.D.		
Labial, to Anterior Cranial Base ..	2.65	2.55	1.6	3.9	1.05	1.06
Downward, from						
Anterior Cranial Base	5.45	2.35	10.0	3.3	4.55	3.41
Edge upward from Mandibular Plane	2.65	1.65	2.0	1.9	0.65	1.23
Edge Labial, to Mandibular Plane ..	1.0	1.45	1.8	1.8	0.8	1.63
Tip (degrees)	3.2	4.65	5.7	5.2	2.5	1.69

** Significant at 0.01 level of confidence.

TABLE 12
Treatment vs. Control
Changes in Vertical Position of Maxilla

	Control		Treatment		dX	t
	Mean	S.D.	Mean	S.D.		
Point A to Anterior Cranial Base ..	4.8	1.65	5.9	2.3	1.1	1.83

TABLE 13
Cervical vs. Control
Changes in Vertical Position of Maxilla

	Control		KC		dX	t
	Mean	S.D.	Mean	S.D.		
Point A to Anterior Cranial Base ..	4.8	1.65	5.8	3.2	1.1	1.3

TABLE 14
Treatment vs. Cervical
Changes in Vertical Position of Maxilla

	Treatment		KC		dX	t
	Mean	S.D.	Mean	S.D.		
Point A to Anterior Cranial Base ..	5.9	2.3	5.8	3.2	0.1	0.12

TABLE 15
Treatment vs. Control
Changes in Horizontal Position of Maxilla

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
Point A to Anterior Cranial Base ..	2.5	1.45	0.8	3.0	3.3	4.65**
Point A to Palatal Plane	0.0	0.65	0.2	1.2	0.2	0.71

** Significant at 0.01 level of confidence.

TABLE 16
Control vs. Cervical
Changes in Horizontal Position of Maxilla

	<i>Control</i>		<i>KC</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
Point A to Anterior Cranial Base ..	2.5	1.45	0.0	2.1	2.5	4.63**
Point A to Palatal Plane	0.0	0.65	0.0	0.5	0.0	—

** Significant at 0.01 level of confidence.

TABLE 17
Treatment vs. Cervical
Changes in Horizontal Position of Maxilla

	<i>Treatment</i>		<i>KC</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
Point A to Anterior Cranial Base ..	-0.8	3.0	0.0	2.1	0.8	1.03
Point A to Palatal Plane	0.2	1.2	0.0	0.5	0.2	0.71

TABLE 18
Treatment vs. Control
Changes in Horizontal Position of Mandible

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
Point B to Anterior Cranial Base ..	1.9	2.55	0.0	3.9	1.9	1.91
Labial						
Point B to S-N at Sella	2.15	3.5	0.3	3.3	1.85	1.81
Labial						
Point B to S-N at Nasion	-1.2	2.85	3.4	3.0	2.2	2.5*
Lingual						
Point B to Mandibular Plane	-0.2	0.35	0.3	0.6	0.1	0.67
Lingual						

Significant at 0.05 level of confidence.

TABLE 19
Treatment vs. Control
Changes in Vertical Position of Mandible

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
Point B to Anterior Cranial Base .. Down	6.15	2.35	9.9	3.0	3.75	4.63**
Point B to S-N Down	6.1	2.7	10.0	3.1	3.9	5.13**
Point B to Mandibular Plane Up	1.35	1.0	1.7	1.7	0.35	0.83

** Significant at 0.01 level of confidence.

TABLE 20
Treatment vs. Control
Changes in Measured Variables

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
Occlusal Plane Angle	1.0	4.1	-0.9	3.0	1.9	1.76
Overbite	0.9	2.15	-1.1	2.6	2.0	2.78*
Overjet	0.1	2.10	-5.6	2.4	5.7	8.38**
Mandibular Plane Angle	1.15	1.4	0.5	1.8	1.65	3.43**
Y-Axis	0.35	1.3	1.9	1.8	1.55	3.30**
Go-Gn/S-N	-1.15	1.90	0.3	1.3	1.45	2.96**
Pogonion to Anterior Cranial Base .. Labial	2.6	2.9	0.1	4.1	2.5	2.36*
Pogonion to Anterior Cranial Base .. Down	7.85	2.1	11.5	3.5	3.65	4.2**

* Significant at 0.05 level of confidence.

** Significant at 0.01 level of confidence.

TABLE 21
Treatment vs. Control
Changes in Face Height

	<i>Control</i>		<i>Treatment</i>		<i>dX</i>	<i>t</i>
	<i>Mean</i>	<i>S.D.</i>	<i>Mean</i>	<i>S.D.</i>		
Upper FH	4.55	1.75	6.1	1.8	1.55	2.92
Lower FH	3.8	1.80	5.5	2.8	1.7	3.24
% Upper FH	0.6	1.2	0.7	0.9	0.1	0.31
% Lower FH	-0.6	1.2	-0.7	0.9	0.1	0.31
Total FH	8.35	2.4	11.7	4.2	3.35	3.25

SUMMARY AND CONCLUSIONS

The quantitative changes which occurred as the result of this specific regimen of treatment in a sample of Class II malocclusions in the early mixed dentition were compared with like changes in a closely matched sample of untreated Class II malocclusions. Selected parameters were also compared with a comparable group of Class II malocclusions treated with only a Kloehe cervical appliance.

1. Treatment as described does tend to slightly increase the vertical eruption of the upper molar.
2. No maxillary orthopedic effects were noted in the horizontal plane.
3. There was significant inhibition of

the forward displacement of the upper molar.

4. Cervical traction without anterior bands resulted in actual distal movement of the upper molar, not just an inhibition of its forward growth.
5. The treatment significantly inhibited the forward growth of the upper incisors and point A.
6. Cervical traction alone had less effect on overjet and the S-N-A angle.
7. A lingual arch is effective in maintaining lower leeway space while still allowing horizontal and vertical growth changes in the positions of molars and incisors.
8. Rotation of the mandible was downward and backward.
9. Overall face height was increased.

REFERENCES

1. Mathews, J. R.: Clinical management and supportive rationale in early orthodontic therapy, *Angle Orthod.*, 31:35-52, 1961.
2. Murray, R. B.: Treatment planning and therapy in the mixed dentition, *Am. J. Orthod.*, 49:641-657, 1963.
3. Dewel, B. F.: Objective of mixed dentition treatment in orthodontics, *Am. J. Orthod.*, 50:504-520, 1964.
4. Funk, A. C.: Mandibular response to headgear therapy and its clinical significance, *Am. J. Orthod.* 53:182-216, 1967.
5. Williams, David R.: The Borderline patient and conservative treatment in the late mixed dentition, *Am. J. Orthod.* 71: 127-155, 1977.
6. Wagers, L. E.: Preorthodontic Guidance and the corrective mixed dentition treatment concept, *Am. J. Orthod.*, 69:1-28, 1976.
7. West, Eugene E.: Treatment objectives in the deciduous dentition, *Am. J. Orthod.*, 55:617, 1969.
8. Watson, Wayne G.: Editorial, First stage treatment-time, methods, potential, *Am. J. Orthod.*, 75:576-579, 1979.
9. Tweed, C. H.: The diagnostic facial triangle in the control of treatment objectives, *Am. J. Orthod.*, 55:651-667, 1969.
10. Meikle, M. C.: The dentomaxillary complex and overjet correction in Class II, division 1 malocclusions, objectives of skeletal and alveolar remodeling, *Am. J. Orthod.*, 77:184-197, 1980.
11. Nagamoto, K. and Yudelson, R.: Early Class II mixed dentition treatment with the universal appliance, *Am. J. Orthod.*, 72:653-664, 1977.
12. Ruff, Roberto M.: Orthodontic treatment in the mixed dentition, *Am. J. Orthod.*, 57:502-518, 1970.
13. Kloehe, S. J.: A new approach to analysis and treatment in the mixed dentition, *Am. J. Orthod.*, 39:161-186, 1953.
14. Weber, Faustin N.: Corrective measures during the mixed dentition, *Am. J. Orthod.*, 43:639-660, 1957.
15. Mathews, J. R.: Interception of Class II malocclusions, *Angle Orthod.*, 41:81-99, 1971.
16. King, E. W.: The possibilities and limitations of mixed dentition treatment, *Am. J. Orthod.* 73:479-498, 1977.

17. Wieslander, L.: Early or late cervical traction therapy of Class II malocclusion in the mixed dentition, *Am. J. Orthod.* 67:423-439, 1975.
18. Kloehn, S. J.: Guiding alveolar growth and eruption of teeth to reduce treatment time and produce a more balanced denture and face, *Angle Orthod.*, 17:10-33, 1947.
19. Kloehn, S. J.: Orthodontics—Force or Persuasion, *Angle Orthod.*, 23:56-66, 1953.
20. Klein, P. L.: An evaluation of cervical traction on the maxilla of the upper first permanent molar, *Angle Orthod.*, 27:61-68, 1957.
21. King, E. W.: Cervical anchorage in Class II, division 1 treatment: A cephalometric appraisal, *Angle Orthod.*, 27:98-104, 1957.
22. Ricketts, R. M.: The influence of orthodontic treatment on facial growth and its development, *Angle Orthod.*, 30:103-133, 1960.
23. Schudy, F. F. and Schudy, G. F.: The Bimetric System, *Am. J. Orthod.* 67:57-80, 1975.
24. Wieslander, L.: The effect of orthodontic treatment on the concurrent development of the craniofacial complex, *Am. J. Orthod.*, 49:15-27, 1963.
25. Schudy, F. F.: Cant of the occlusal plane and axial inclination of teeth, *Angle Orthod.*, 33:69-82, 1963.
26. Newcomb, M. R.: Some observations on extraoral treatment, *Angle Orthod.*, 28:131-148, 1958.
27. Poulton, D. R.: The influence of extraoral traction, *Am. J. Orthod.*, 53:8-18, 1967.
28. Poulton, D. R.: Changes in Class II malocclusions with and without occipital headgear therapy, *Angle Orthod.*, 29:234-250, 1959.
29. Schudy, F. F.: The rotations of the mandible resulting from growth: Its implications in orthodontic treatment, *Angle Orthod.*, 35:36-50, 1965.
30. Root, T. L.: Interview on headgear, *J. Clin. Orthod.*, 9:20-41, 1975.
31. Thurow, R. C.: Craniomaxillary orthopedic correction with en masse dental control, *Am. J. Orthod.*, 68:601-624, 1975.
32. Schudy, F. F.: Vertical growth versus anteroposterior growth as related to function and treatment, *Angle Orthod.*, 34:75-93, 1964.
33. Merrifield, L. and Cross, J. J.: Directional forces, *Am. J. Orthod.*, 57:435-464, 1970.
34. Mills, C. M., Holman, R. G., and Graber, T. M.: Heavy intermittent cervical traction in Class II treatment: A longitudinal cephalometric assessment, *Am. J. Orthod.*, 74:361-379, 1978.
35. Wieslander, L.: The effect of force on craniofacial development, *Am. J. Orthod.*, 65:531-538, 1974.
36. Wieslander, L., and Buck, D. L.: Physiologic recovery after headgear treatment, *Am. J. Orthod.*, 66:294-301, 1974.
37. Ringenberg, Q. M. and Butts, W. C.: A controlled cephalometric evaluation of single arch cervical traction therapy, *Am. J. Orthod.*, 57:179-185, 1970.
38. Watson, W. G.: A computerized appraisal of the high-pull face bow, *Am. J. Orthod.*, 62:561-579, 1972.
39. Melson, B.: Effects of cervical anchorage during and after treatment: An implant study, *Am. J. Orthod.*, 73:526-540, 1978.
40. Baumrind, S., Molthen, R., West, E. E., and Miller, D. M.: Mandibular plane changes during maxillary retraction, *Am. J. Orthod.*, 74:32-40, 1978.
41. Baumrind, S., Molthen, R., West, E. E., and Miller, D. M.: Part 2 Mandibular plane changes during maxillary retraction, *Am. J. Orthod.*, 74:603-620, 1978.
- Addenda, *Am. J. Orthod.*, 75:86-89, 1979.
42. Baumrind, S., Molthen, R., West, E. E., and Miller, D. M.: Distal displacement of the maxilla and upper first molar, *Am. J. Orthod.*, 75:630-640, 1979.
43. Graber, T. M., Chung, D. D. B., and Aoba, J. T. L.: Dentofacial orthopedics versus orthodontics, *J. Am. Dent. Assoc.*, 75:1145-1166, 1967.
44. Singer, J.: The effect of the passive lingual archwire on the lower denture, *Angle Orthod.*, 44:146-155, 1974.
45. Moyers, R. E. and Bookstein, F. L.: The inappropriateness of conventional cephalometrics, *Am. J. Orthod.*, 75:599-617, 1979.
46. Baumrind, S. and Miller, D. M.: Computer-aided headfilm analysis: the University of California, San Francisco, in publication.