

An orthodontic study of temporomandibular joint disorders

Part 1: Epidemiological research in Japanese 6-18 year olds

Etsuko Moteji, DDS, DDSc; Haruyo Miyazaki, DDS; Isao Ogura, DDS; Harumi Konishi, DDS; and Masayuki Sebata, DDS, DMSc

In recent years, an increasing number of patients have complained of temporomandibular joint disorders (TMD).^{1,2} In an attempt to identify the causes of TMD, the type of occlusion in children with TMD symptoms was recorded.

There have been a number of epidemiologic investigations of children with TMD symptoms³⁻²⁰ (Table 1). Since approaches and methodologies differ, the prevalence varies in these studies from 10% to 80%. Interpreting these findings requires careful consideration of the peculiarities of each investigation.^{6,21}

Materials and methods

This study included 7337 children (Table 2): 3219 boys and 4118 girls from four elementary schools, four junior high schools and three high schools in Chiba and Tokyo, Japan. The data was collected from April 1986 through June 1986 by oral examination during regularly held physical examinations.

First we taught the subjects to locate their TM joint, then we asked them about past or recent TMJ sound, pain or abnormal jaw movements. Then, while palpating the TMJ, we had the subjects open and close their mouth several times to confirm the

Abstract

Malocclusion is considered one of the etiological factors of temporomandibular joint disorder (TMD). The purpose of this study was to investigate the prevalence of TMD and the relationship between TMD and the type of occlusion. The sample consisted of 7337 Japanese children, 6-18 years old, 3219 boys and 4118 girls. TMD symptoms were recorded as well as the type of occlusion in children with TMD.

The prevalence of TMD overall was 12.2%. The prevalence increased with age and was slightly higher in girls (13%) than in boys 11.1%. This difference was not statistically significant. Joint sound as the only symptom was more common in younger subjects. TMD symptoms seemed more complicated with age when pain and abnormal jaw movement combined with sound. Joint sound was the most common symptom (89.3%), followed by the combination of sound and pain (2.2%). The incidence of other symptoms was under 1%. In subjects with TMD, 24.9% exhibited crowding, 20.1% had excessive overjet, 6.8% deep bite, 6.3% edge-to-edge bite, 5.6% anterior crossbite, 5.4% open bite, and 3.8% posterior crossbite. Morphologically normal occlusion was observed in 27.1%. In this study, many subjects with TMD had malocclusions. Early treatment may be important in the prevention of severe TMD. Although those with morphologically normal occlusions were included, a more detailed study concerning other causes of TMD is needed also.

This manuscript was submitted February 1992. It was revised and accepted for publication June 1992.

Key Words

Malocclusion • Temporomandibular joint disorder • Epidemiological research

Table 1
Epidemiological and clinical studies on children with TMD

Researcher	Number of Subjects	Male:Female	Age	Prevalence symptom /sign	
Geering-Gaerney & Rakosi (71)	241	—	8-14	—	41%
Agerberg& Carlsson (72)	1106	531:575	15-74	57%	—
Ingerval & Hedegard (74)	287	287:	18-20	12%	—
Helkimo (74)	321	156:165	15-65	57%	88%
Siebert(75)	98	—	12-16	—	80%
Molin et al (76)	253	253:	18-25	12%	28%
Dibbets (77)	112	49:63	8-17	—	46%
Grosfeld & Czarnecka (77)	250	114:136	6-8	—	56%
W-Markowerowa et al (77)	2100	117:133	13-15	—	68%
Williamson (77)	304	129:175	10-15	—	40%+Brux 27%+Brux
Helkimo et al (79)	58	0:58	6-16	—	35%
E-Eriksson et al (81)	136	62:74	18-28	45%	60%
	131	70:61	7	16%	33%
	135	76:59	11	17%	46%
			15	25%	61%
Magnusson, E-Eriksson, et al. (85)	119	59:61	11	62% (11%mod.)	
			15	66% (17%mod.)	
Ogura, et al (85)	2198	1095:1103	10-18	9.8%	
Ishibashi,et al. (85)	1113	561:466	12-17	30.7%	
Grosfeld, et al (85)	400	197:203	15-18	—	68.25%
	400	208:192	19-22	—	67%
Brandt (85)	1342	673:669	6-17	10.74%	
Nilner (86)	440	222:218	7-14	36%	39%
	309	147:162	15-18	41%	34%
Bernal (86)	149	79:70	3-5		21.1%
Wanman, et al (86)	285	146:139	17	62%	
Szentpetery (87)	85	56:29	11-20	16.5%	67.1%
Motegi, et al (88)	7337	3219:4118	6-18	12.2%	

Table 2
Number of Subjects

Grade	Age	Male	Female	Total
E	1 6	198	197	395
	2 7	207	209	416
	3 8	222	201	423
	4 9	224	226	450
	5 10	239	258	497
	6 11	289	265	554
J	1 12	397	395	792
	2 13	374	370	744
	3 14	375	330	705
H	1 15	217	544	761
	2 16	215	542	757
	3 17	262	582	846
E Total		1379	1356	2735
J Total		1146	1092	2238
H Total		694	1670	2346
Total		3219	4118	7337

Table 3
Prevalence of TMD by age and sex

Grade	Age	Male		Female		Total	
		Number	%	Number	%	Number	%
E	1 6	5	2.5	11	5.6	16	4.1
	2 7	7	3.4	11	5.3	18	4.3
	3 8	15	6.8	14	7.0	29	6.9
	4 9	9	4.0	10	4.4	19	4.2
	5 10	16	6.7	21	8.1	37	7.4
	6 11	15	5.2	22	8.3	37	6.7
J	1 12	51	12.9	48	12.2	99	12.5
	2 13	59	15.8	41	11.2	100	13.5
	3 14	58	15.5	68	20.6	126	17.9
H	1 15	41	18.9	94	17.3	135	17.7
	2 16	35	16.3	93	17.2	128	16.9
	3 17	46	17.6	103	17.6	149	17.6
E Total		67	4.9	89	6.6	156	5.7
J Total		168	14.7	157	14.4	325	14.5
H Total		122	17.6	290	17.4	412	17.4
Total		357	11.1	536	13.0	893	12.2

findings derived by questioning the subjects. Finally, we observed and classified the occlusions of those with TMD and asked whether or not they had received orthodontic or any other treatment for TMD.

Classification of symptoms

Sound: clicking or crepitus

Pain: pain that occurs from opening and closing the mouth, pain from chewing food and any pain in the TMJ

Abnormal jaw movement: less than two fingers of childrens' own in the open mouth, and difficulty opening and closing the mouth.

Classification of occlusion

Excessive overjet: overjet exceeding 5 mm

Anterior crossbite: at least 4 anterior teeth in crossbite

Edge-to-edge bite: both overbite and overjet 0mm

Crowding: severe crowding with normal relationship of the maxilla to the mandible

Open bite: negative overbite

Deep bite: overbite exceeding 5mm with small overjet

Posterior crossbite: molar teeth that exhibit crossbite

Morphologically normal occlusion: no abnormality in tooth alignment.

Using these standards, the research was conducted by four dentists known to the authors.

Results

The prevalence of TMD symptoms for all ages was 12.2%. The elementary school level was 5.7%,

the junior high school level was 14.5% and the high school level was 17.4% (Table 3). These findings show a gradual increase in prevalence with age. School grade differences were significant respectively at the 1% level according to Chi-square analysis.

The prevalence of TMD symptoms was 11.0% for boys and 13.0% for girls. However, the sex difference was not significant at the 1% level according to the Chi-square analysis.

We used seven categories of symptoms: sound, pain, abnormal jaw movement and four combinations of these three. The most common symptom at the elementary school level was sound. The number of children with sound as a symptom increased at the junior high and high school levels. Combined problems started appearing at the junior high and high school levels. (Table 4)

The most common symptom was sound (97.2%), followed by pain (10.2%) and abnormal jaw movement (0.9%) (Table 5). Sound alone accounted for 89.3% of the symptoms, the combination of sound and pain accounted for 7.6%, pain alone 2.2% and other symptoms less than 1.00%.

There was a high percentage of TMD in cases of crowding and excessive overjet and the prevalence of each increased with age (Table 6). Morphologically normal occlusion peaked around the transition period from elementary school to junior high school. Anterior crossbite, edge-to-edge bite, open bite, deep bite and posterior crossbite exhibited a very small prevalence of symptoms and showed little increase with age.

E: Elementary school
J: Junior high school
H: High school

E: Elementary school
J: Junior high school
H: High school

S: Sound
P: Pain
A: Abnormal jaw movement

Table 4
Age difference by symptom

	Grade	Age	S	S+P	P	S+A	S+P+A	A	P+A	Total
E	1	6	3.8	0.3	0	0	0	0	0	4.1
	2	7	4.1	0	0.2	0	0	0	0	4.3
	3	8	6.6	0	0.3	0	0	0	0	6.9
	4	9	4.2	0	0	0	0	0	0	4.2
	5	10	7.0	0	0.4	0	0	0	0	7.4
	6	11	6.7	0	0	0	0	0	0	6.7
J	1	12	10.2	1.8	0.4	0	0.1	0	0	12.5
	2	13	12.0	0.7	0.4	0.1	0.1	0.1	0	13.5
	3	14	15.5	1.6	0.6	0.1	0.1	0	0	17.9
H	1	15	16.3	0.9	0.4	0	0	0.1	0	17.7
	2	16	15.2	1.5	0.1	0	0	0	0.1	16.9
	3	17	15.0	2.3	0.2	0.1	0	0	0	17.6
Total			10.8	0.9	0.3	0.1	0.1	0.0	0.0	12.2

Table 5
Percentage of each symptom

Symptoms	Number	%
S	796	89.3
S+P	68	7.6
P	20	2.2
S+A	3	0.3
S+P+A	3	0.3
A	2	0.2
P+A	1	0.1
Total	893	100.0

The percentages of various types of occlusion among those with TMD were as follows: crowding, 24.9%; excessive overjet, 20.1%; deep bite, 6.8%; edge-to-edge bite, 6.3%; anterior crossbite, 5.6%; open bite, 5.4%; posterior crossbite, 3.8% and morphologically normal occlusion, 27.1%.

Twelve children had received some kind of treatment for TMD. None were elementary school students, 0.6% were junior high and 2.4% were high school students. Five children had received orthodontic treatment.

Discussion

In general, our questions to the children regarding TMD were not usually well understood if they had never recognized such problems before. For this reason, we had them locate their TMJ by lightly

touching the front area of their tragus. We then asked whether or not they had a current and/or past history in regard to TMD. Through palpation we further confirmed the findings gathered from the questions. Differentiating joint pain, toothache and earache seemed very difficult.

Some dentists insist on the importance of auscultation with a stethoscope, but we believe palpation provides a more reliable method for recognizing TMJ sound. There is a difficulty in examination in such a noisy classroom using a stethoscope. The standard method should be simple and reproducible in epidemiological research of this type.⁶

As mentioned above, the occlusal condition of those with TMD was examined. Only a few of the subjects were aware of symptoms. Some children said they had created sound or purposely dislocated their jaws for fun, and were simply unaware of having a temporomandibular disorder.

The examination was based on three main symptoms in order to simplify the classification. The categories selected included the identification of a single symptom and combined symptoms^{16,17} in order to make the analysis of the level of severity less complicated.

The classification of occlusion should include any abnormality of upper and lower, left and right, anterior and posterior parts.

The prevalence of TMD in Japan is rather low compared with other countries. Each researcher samples different populations: e.g. elderly subjects or orthodontic patients. There is also a difference in information gained through questionnaires which

N: Morphologically normal occlusion
 C: Crowding
 EJ: Excessive overjet
 D: Deep bite
 EE: Edge-to-edge bite
 A: Anterior crossbite
 O: Open bite
 PC: Posterior crossbite

Table 6
Age difference in the prevalence of TMD symptoms

Grade	Age	N	C	EJ	D	EE	A	O	P	Total	(%)
E	1 6	1.3	0.3	0	0	0	1.0	1.5	0	0	4.1
	2 7	1.2	0.7	0.5	0.5	0.2	0.7	0.5	0	0	4.3
	3 8	2.4	1.2	1.4	0.5	0.2	0.5	0.5	0.2	0.2	6.9
	4 9	0.7	0.6	1.3	0.2	0.2	0.7	0.2	0.2	0.2	4.2
	5 10	2.0	1.2	2.4	0.4	0.2	0.6	0.2	0.2	0.4	7.4
	6 11	1.1	2.5	2.3	0.2	0.2	0.4	0	0	0	6.7
J	1 12	4.9	3.0	1.5	1.8	0.4	0.8	0	0	0.3	12.5
	2 13	6.0	2.3	1.7	0.9	0.8	0.1	0.9	0.4	0.4	13.5
	3 14	6.5	3.6	2.8	1.6	1.0	0.9	0.7	0.9	0.9	17.9
H	1 15	3.7	4.8	3.8	0.9	2.0	0.7	1.1	0.9	0.9	17.7
	2 16	2.6	4.6	3.8	1.2	1.8	1.3	0.8	0.7	0.7	16.9
	3 17	3.0	6.4	4.5	0.5	0.7	0.6	1.2	0.8	0.8	17.6
Total		3.3	3.0	2.5	0.8	0.8	0.7	0.7	0.5	0.5	12.2

depend only on people's perceptions, and information gained by doctors, especially in children who may not be aware of their condition.

This study showed an increasing trend in TMD with age similar to the Scammon's mandibular growth curve. From this finding, we believe it is important to prevent TMD at an early age.

Although several clinical studies²²⁻²⁶ have shown more women have TMD than men, our research found no differences by sex. Other epidemiological reports^{16,17,27} have also shown no difference between the sexes. We believe the sex difference could be due to the increased frequency of medical visits by women.

Sound was the main symptom identified at the elementary school level. After the first year of junior high school, the increase in sound was significant. The prevalence of other symptoms was very low, but definitely showed a tendency to increase. Also, combinations of symptoms increased with age. These findings indicate the need for diagnosis and treatment at an early age.¹

Only 12 children had seen doctors to complain of TMD as their main reason to visit (0.1% of the whole). Of those 12 children, 10 had sound and pain, one had pain alone and one had abnormal jaw movement as the main complaint for the visit; none of them went to see a doctor because of sound alone. Although the numbers are small, it appears that people do not see a doctor for sound unless it is accompanied by pain and/or abnormal jaw movement.²²

The incidence of excessive overjet and crowding


Table 7
Prevalence by type of occlusion

Type of occlusion	Number	%
Morphologically normal occlusion	242	27.1
Crowding	222	24.9
Excessive overjet	180	20.1
Deep bite	61	6.8
Edge-to-edge bite	56	6.3
Anterior crossbite	50	5.6
Open bite	48	5.4
Posterior crossbite	34	3.8
Total	893	100.0

increased with age. With excessive overjet, people create a habit of protruding the mandible.²⁸ This tends to cause a dual bite^{29,30} which could, over time, aggravate the muscles of mastication and muscle tension and load the TM joints. Crowding tends to cause occlusal interferences and seems to be a critical factor in TMD.¹¹⁻¹⁴

Anterior crossbite, skeletal especially, is less likely to interfere with occlusion during chewing.^{31,32} If it creates an occlusal interference, it seems to occur most commonly with the anterior teeth and hardly seems to affect the TMJ or nearby muscles.³¹ However, Mukaizawa reported that patients with an anterior crossbite are most likely to develop the symptoms of TMD at an early age. This finding demands further research.

Figure 1
Age difference in the prevalence of the TMD by occlusion N: Morphologically normal occlusion C: Crowding EJ: Excessive overjet D: Deep bite EE: Edge-to-edge bite AA: Anterior crossbite O: Open bite PC: Posterior crossbite



An edge-to-edge bite has little interference and a relatively low prevalence in TMD. From this we can hypothesize that TMD may be related to the longstanding presence of occlusal interferences.

The prevalences of open bite, posterior crossbite and deep bite were small and we found no significant increase in symptoms with age. However, from clinical experience, we think TMD may accompany open bite, which is an unstable mandibular position, deep bite, which tends to push the mandible posteriorly, and posterior crossbite, which forces the mandibular position either to the left or the right. The incidence of morphologically normal occlusion with TMD reaches its peak in the upper grade levels of elementary and junior high school. This is largely related to the drastic increase in the prevalence of TMD at this age. The increase may be attributed to stress, such as from entrance examinations, or inflammation caused by the eruption of second molars.

Nilner¹¹⁻¹⁴ reported that the percentage of children

with deep bite is high, while Mineno³³ noted that excessive overjet and crowding are common among TMD patients. Williamson³⁴ reported Angle Class I and Class II relationships were the most common occlusal types. Our findings by occlusal type show morphologically normal occlusion, crowding and excessive overjet to be most frequent with others making up a relatively small percentage.

Some reports^{2,29,30} on malocclusion are from a functional point of view. Likewise, each malocclusion should be classified based on function. The fact that a large percentage of children have a morphologically normal occlusion indicates that an abnormal a-p relationship or malalignment is not always a major cause of TMD; other problems such as occlusal interference, abnormality in function and formation of the TMJ, trauma, parafunction, and habitual unilateral chewing, which are not all classified under the type of occlusion, could be causes of TMD.

Of the 12 children who had received treatment for

their TMD, six had first visited orthopedic surgeons, five had visited dentists, and one had seen a chiropractor. This suggests that dentists could become more widely recognized as providers of TMD treatment. We believe people need to be educated about this.^{21,35}

Of those who had TMD, only five had received orthodontic treatment; therefore, the study of Part 2 will follow this.

Conclusions

The incidence of TMD increases with age in children and adolescents and its symptoms tend to become more complex with age. Most children with TMD have an abnormal occlusion, indicating a need for further and more detailed investigation into the pathogenesis of TMD.

Acknowledgments

The present study was supported by Grant-in-Aid for Scientific Research (C) 63570972.

The author would like to thank Dr. H.T. Perry for suggesting and encouraging publication of this study in English

Author Address

Dr. Etsuko Motegi
Department of Orthodontics
Tokyo Dental College
1-2-2 Masago, Mihama-ku
Chiba-shi, Chiba-ken
261, Japan

E. Motegi is Assistant Professor in the Department of Orthodontics, Tokyo Dental College.

H. Miyazaki is Assistant Professor in the Department of Orthodontics, Tokyo Dental College.

I. Ogura is in private practice in Chiba-shi, Chiba-ken.

H. Konishi is in private practice in Ichihara-shi, Chiba-ken.

M. Sebata is chairman and Professor, Department of Orthodontics, Tokyo Dental College.

References

- 1). Mukaiyama T, Fukazawa H, et al. Prevalence of temporomandibular dysfunction with anterior crossbite malocclusion in 6-10 years old Japanese children. *J Japan Ortho Society* 1986;45:574-80.
- 2). Nohmi Y, Ohtsuiji T, et al. Occlusal features of orthodontic patients with symptoms of temporomandibular joint dysfunction. *J Japan Ortho Society* 1987;46:696-707.
- 3). Hansson T and Nilner M. A study of the occurrence of symptoms of diseases of the temporomandibular joint masticatory musculature and related structures. *J Oral Rehab* 1975;2:313-324.
- 4). Grosfeld O and Czarnecka B. Musculoarticular disorders of the stomatognathic system in school children examined according to clinical criteria. *J Oral Rehab* 1977;4:193-200.
- 5). Grosfeld O, Jackowska M and Czarnecka B. Results of epidemiological examinations of the temporomandibular joint in adolescents and young adults. *J Oral Rehab* 1985;12:95-105.
- 6). Helkimo MI. Epidemiological surveys of dysfunction of the masticatory system. In *Temporomandibular Joint Function and Dysfunction*, eds Zarb GA and Carlsson GE, Munksgaard, Copenhagen, 1979, The CV Mosby Company, St. Louis, Missouri, 175-192.
- 7). Wigdorowicz-Makowerowa N, Grodzki C, et al. Epidemiologic studies on prevalence and etiology of functional disturbances of the masticatory system. *J Prosthet Dent* 1979;41:76-82.
- 8). Egermark-Eriksson I, Carlsson, Inger-vall B. Prevalence of mandibular dysfunction and orofacial parafunction in 7, 11 and 15-year-old Swedish children. *Eur J Orthod* 1981;3:163-172.
- 9). Egermark-Eriksson I, Ingervall B. Anomalies of occlusion predisposing to occlusal interference in children. *Angle Orthodont* 1982;52:293-299.
- 10). Egermark-Eriksson I, Ingervall B and Carlsson GE. The dependence of mandibular dysfunction in children on functional and morphologic malocclusion. *Am J Orthod* 1983;83:187-194.
- 11). Nilner M. Prevalence of functional disturbances and diseases of the stomatognathic system in 7-14 year olds. *Swed Dent J* 1981;5:173-187.
- 12). Nilner M. Prevalence of functional disturbances and diseases of the stomatognathic system in 15-18 year olds. *Swed Dent J* 1981;5:189-197.
- 13). Nilner M. Functional disturbances and diseases in the stomatognathic system among 7-to 18-year olds. *J Cranio Practice* 1985;3:358-367.
- 14). Nilner M. Functional disturbances and diseases of

- the stomatognathic system. A cross-sectional study. *J Pedodont* 1986;10:211-238.
- 15). Magnusson T, Egermark-Eriksson I and Carlsson GE. Four year longitudinal study of mandibular dysfunction in children. *Comm Dent Oral Epidemiol* 1985;13:117-120.
- 16). Ogura T. Prevalence and distribution of the symptoms of TMJ dysfunction syndrome in adolescents. *Japan Dent Review* 1985;512:149.
- 17). Ohno H, Morinushi T, et al. Prevalence and distribution of the symptoms of TMJ dysfunction syndrome with adolescents. *Japan J Ped* 1985;23:94-102.
- 18). Wanman A and Agermark G. Relationship between signs and symptoms of mandibular dysfunction in adolescents. *Comm Dent Oral Epidemiol* 1986;14:225-230.
- 19). Brandt D. Temporomandibular disorders and their association with morphologic malocclusion in children. Development aspects of temporomandibular joint disorders. Center for Human Growth and Development. The University of Michigan, Michigan, 1985.
- 20). Szentpetery A, Fazekas A and Mari A. An epidemiologic study of mandibular dysfunction dependence on different variable. *Comm Dent Oral Epidemiol* 1987;15:164-168.
- 21). Kent G. Prevalence vs. incidence of the mandibular pain dysfunction syndrome: Implications for epidemiological research. *Comm Dent Oral Epidemiol* 1985;13:113-116.
- 22). Fujita H, Kanai Y, et al. Clinicostatical study of temporomandibular joint pain-dysfunction-syndrome. *J Japan Oral Surg* 1980;26:1508-15.
- 23). Furuta H, Kuwahara M, et al. Clinical study on TMJ arthrosis in young people. Discussion on the dentocraniofacial morphology. *J Japan Oral Surg* 1987;33:257-63.
- 24). Taguchi N, Maruyama T, et al. Clinicostatistical study of temporomandibular arthrosis. *J Japan Oral Surg* 1986;32:399-405.
- 25). Taguchi N, Kuwahara M, et al. Clinical study of temporomandibular arthrosis. *J Japan Oral Science* 1986;35:46-60.
- 26). Takada K, Fukuda K, et al. Clinical study of temporomandibular joint disturbance. *J Osaka Univ* 1968;13:291-95.
- 27). Solberg WK, Woo MW and Houston JB. Prevalence of mandibular dysfunction in young adults. *JADA* 1979;98:25-34.
- 28). Snow DF. Diseases of the temporomandibular apparatus. 2nd ed. St. Louis, The CV Mosby Company 1982;83.
- 29). Ricketts RM. A study of changes in temporomandibular relations associated with the treatment of Class II malocclusion. *Am J Orthod* 1952;38:918-933.
- 30). Ricketts RM. Abnormal function of the temporomandibular joint. *Am J Orthod* 1955;41:435-441.
- 31). Nakazawa K. Analysis of mandibular movement in the patients with temporomandibular joint arthrosis. *Shikwa Gakuho* 1974;74:277-311.
- 32). Shibata T. Studies on mandibular movements in the patients with mandibular protrusion. *Shikwa Gakuho* 1981;81:241-265.
- 33). Mineno Y, Taguchi N, et al. Odontological studies on temporomandibular joint arthrosis in young people. Studies on occlusion and tooth material. *J Japan Oral Surg* 1986;32:1908-16.
- 34). Williamson EH. Temporomandibular dysfunction in pretreatment adolescent patients. *Am J Orthod* 1977;72:429-433.
- 35). Akamine E, Takenoshita Y, et al. Clinicostatistical observations the temporomandibular arthrosis. *J Japan Oral Surg* 1977;23:243-49.