

Vertical Growth Changes After Adenoidectomy

Tülin Arun, DDS, PhD^a; Fulya Isik, DDS, PhD^b; Korkmaz Sayinsu, DDS, PhD^b

Abstract: The purpose of this retrospective investigation is to compare vertical growth component of craniofacial structure of subjects with early and late adenoidectomy history. The study consisted of 93 lateral cephalometric radiographs of three groups of randomly selected patients. The first group was made up of 12 patients (10 male and two female) with an average age of 11.16 ± 2.08 years, who had been operated upon between 1.5–4 years of age. The second group was made up of 54 patients (25 male and 29 female) with an average age of 12.18 ± 2.6 years, who had been operated upon after four years of age. The third group of 27 patients (7 male and 20 female) with clear airway with an average age of 11.18 ± 2.35 years was used as the control. The data obtained from two adenoidectomy groups were compared and because no statistically significant difference was found except for ANSMe/NMe, the two groups were pooled and compared with the growth pattern in the control sample. There were statistically significant differences in the following parameters: SNGoMe, PPGoMe angle, Gonial angle, Gonial ratio, Σ of inner angles, ANSMe/NMe ratio, Jarabak ratio, PNS-ad1 distance, PNS-ad2 distance, OAW1 distance. When compared with the control group, the adenoidectomy group showed a more vertically directed growth pattern, however, there were no vertical growth pattern differences between the two groups of children who had adenoidectomy before and after four years of age. (*Angle Orthod* 2003;73:146–150.)

Key Words: Adenoidectomy; Growth pattern; Upper airway inadequacy

INTRODUCTION

Because nasal airway inadequacy is one of the major environmental causes of malocclusions observed in growing children, many studies have been carried out to reveal the association between nasal respiratory impairment and dentofacial developmental anomalies.^{1–9} Investigations carried out on the subject have pointed out the role of enlarged adenoids,^{1,10–12} allergic rhinitis^{7,13,14} and enlarged tonsils¹⁵ on the effective mode of breathing.

In a serial study of 120 Canadian boys aged 6–18 years, Woodside and Linder-Aronson¹² found that 26% of the subjects had short anterior face heights throughout their growth periods and 100% of this group had clear pharyngeal airways as assessed from annual serial lateral cephalograms and 90% of the group showed clear nasal passages as assessed from annual serial posteroanterior cephalograms. Also, in the 22 males who had a longer or increased lower face height, the mean size of the airway through the nose

or nasopharynx was narrower than that in the control group. In 1970, Linder-Aronson in another study found that the craniofacial morphology of children with upper airway obstruction because of enlarged adenoids differed systematically from that of a matched control group.¹ An increase in anterior facial height and gonial angle and a larger mandibular plane angle with a reduced facial prognathism were the major characteristics of these adenoid children. Linder-Aronson in a follow-up study,¹⁰ reported that some children may demonstrate varying degrees of spontaneous recovery during the first year after the change from mouth to nasal breathing, which was assessed by the authors clinically after adenoidectomy. These changes were reported to occur during the five-year period after adenoidectomy, although not reaching control values.¹¹

Besides these studies, which concluded that mouth breathing disrupts the muscle forces exerted by the tongue, cheeks, and lips on the dental arches and produce unwanted growth pattern,^{1–3,5,6,10–12,16–18} there are other studies that deny a significant relationship between facial morphology and mode of breathing.^{4,8,19–23} Gwynne-Evans and Ballard,¹⁹ in their 15-year longitudinal study, concluded that orofacial morphology remains constant during growth, regardless of breathing patterns. Similarly, Humphery and Leighton⁴ studied 1033 children and reported an approximately equal distribution of malocclusions in nose and mouth breathers. These authors also noted that of those children who kept their mouths open while breathing, al-

^a Associate Professor, Faculty of Dentistry, Yeditepe University, İstanbul, Turkey.

^b Assistant Professor, Faculty of Dentistry, Teditepe University, Göztepe, İstanbul, Turkey.

Corresponding author: Tülin Arun, Yeditepe University, Faculty of Dentistry, Bağdat cad. No. 238 Göztepe, İstanbul, Turkey. (e-mail: tarun@yeditepe.edu.tr)

Accepted: July 2002. Submitted: May 2002.

© 2003 by The EH Angle Education and Research Foundation, Inc.

TABLE 1. Gender Distribution of Subjects

| | Boys | Girls | Total |
|-------------------------------------------|------|-------|-------|
| Had adenoidectomy between 1.5 and 4 years | 10 | 2 | 12 |
| Had adenoidectomy after four years | 25 | 29 | 54 |
| Without nasal obstruction history | 7 | 20 | 27 |
| Total | 42 | 51 | 93 |

most half respired nasally. Watson et al,⁸ and Rasmus and Jacobs⁹ found no association between rhinomanometric measures of airway adequacy and type of malocclusion or craniofacial morphology.

In long-face patients, the orthodontist's goal is to find an evidence-based explanation of etiology and, if possible, modify the growth direction to create a more favorable craniofacial skeletal morphology. Because the largest increase in growth takes place in the very early years of life, and by the age four years the craniofacial skeleton reaches 60% of its adult size,²⁴ timing of the removal of excessive adenoid tissue should play an important role on the recovery of this undesirable craniofacial growth pattern often termed as long-face syndrome or adenoid face, if there is a relationship between large adenoids and facial morphology.

Cross-sectional studies carried out on this subject are interested mostly in the mode of respiration in persons with selected facial features.²⁵ The purpose of our retrospective cross-sectional investigation is to compare the vertical growth component of craniofacial structure of subjects with early and late adenoidectomy history with controls.

MATERIALS AND METHOD

The study material consisted of a total of 93 lateral cephalometric radiographs of 42 boys and 51 girls, accumulated during the years 2000 to 2001, which were images of patients who were admitted to the University Department of Orthodontics for treatment of various malocclusions.

The radiographs were divided into three groups (Table 1). Group I was made up of the radiographs of 12 patients (10 male and two female; age 11.16 ± 2.08 years) who had an adenoidectomy between 1.5–4 years of age. Group II was composed of the radiographs of 54 patients (25 male and 29 female; age 12.18 ± 2.6 years) who had been operated upon after four years of age. Measurements for the first two groups were obtained from cephalometric radiographs, which were taken at least one year after adenoidectomy in the growing children. Group III, the control, was composed of the radiographs of 27 patients (7 male and 20 female; age 11.18 ± 2.35 years) with a clear airway.

The adenoidectomy group patients had been referred for the surgical procedures because of their obstructed nasal breathing. The control group had no history of nasal obstruction, nasorespiratory allergy or recurrent otitis media. None of the patients had undergone any kind of orthodontic treatment. These subjects were randomly selected from the department archives taking their age and their airway clear-




FIGURE 1. Nasopharyngeal and oropharyngeal airway measurements: 1. PNS-ad1 distance, 2. PNS-ad2 distance, 3. OAW1 distance, 4. OAW2 distance, 5. OAW3 distance.




FIGURE 2. SNGoMe angle, Gonial ratio, ANSMenMe ratio, Jarabek ratio, PPGoMe angle measurements—comparison of adenoidectomy and control groups.

ance—as seen from lateral cephalometric film—into consideration.

Measurements and statistical analysis

For the assessment of vertical growth direction, SNGoMe angle, PPGoMe angle, gonial angle, gonial ratio, Σ (sum




FIGURE 3. Gonial angle and sum of inner angle measurements—comparison of adenoideotomy and control groups.




FIGURE 4. Pharyngeal measurements—comparison of adenoideotomy and control groups.

of the internal cranial angles), ANSM_e/Nme ratio, and Jarabak were measured. The nasopharyngeal and oropharyngeal airways were investigated by PNS-ad1, PNS-ad2, OAW1, OAW2, and OAW3 distances as defined by earlier research^{26–28} (Figure 1).

For the detection of method error, 25 radiographs were randomly selected from the study material and retraced and remeasured one week after the first measurement. Paired *t*-tests gave no errors related to the measurements. The Mann-Whitney *U*-test from the SPSS statistics program was used for analyzing the data.

RESULTS

The data obtained from two adenoideotomy groups were compared and because no statistically significant difference was found except for ANSM_e/Nme ($P < .05$) (Table 2),

TABLE 2. Comparison Between Two Different Groups of Adenoideotomy^a

| | Adenoidectomy | | | | Test (Mann-Whitney <i>U</i>) |
|--------------|-------------------------|------|----------------------|-------|----------------------------------|
| | (1.5–4 years) n = 12 | | (>4 years) n = 54 | | |
| | Mean | SD | Mean | SD | |
| SNGoMe | 41.17 | 6.00 | 40.71 | 6.34 | |
| SNPP | 7.25 | 2.75 | 8.58 | 3.42 | |
| PPGoMe | 33.83 | 5.94 | 31.71 | 6.58 | |
| Gonial angle | 129.50 | 7.97 | 130.74 | 6.03 | |
| Gonial ratio | 66.13 | 5.79 | 63.45 | 7.78 | |
| Σ | 403.17 | 6.84 | 400.76 | 6.82 | |
| ANSMe/Nme | 58.46 | 2.51 | 56.90 | 2.14 | * |
| Jarabak | 59.23 | 3.92 | 60.94 | 4.60 | |
| Sar/Ramus | 85.63 | 7.18 | 83.33 | 10.80 | |
| PNS-ad1 | 22.83 | 5.38 | 21.76 | 5.96 | |
| PNS-ad2 | 17.00 | 5.17 | 17.24 | 5.38 | |
| OAW1 | 9.67 | 3.10 | 9.07 | 2.95 | |
| OAW2 | 9.67 | 3.68 | 10.78 | 4.74 | |
| OAW3 | 12.54 | 2.64 | 12.51 | 3.61 | |

^a * $P < .05$.

TABLE 3. Statistical Evaluation of Adenoideotomy Patients^a

| | Adenoideotomy Group n = 66 | | Control Group n = 27 | | Test (Mann-Whitney <i>U</i>) |
|------------------------|----------------------------|-------|----------------------|-------|-------------------------------|
| | Mean | SD | Mean | SD | |
| SNGoMe | 40.80 | 6.24 | 35.70 | 5.91 | ** |
| SNPP | 8.34 | 3.33 | 8.78 | 3.85 | |
| PPGoMe | 32.10 | 6.47 | 26.93 | 7.14 | ** |
| Gonial angle | 130.52 | 6.37 | 126.87 | 8.32 | * |
| Gonial ratio | 63.94 | 7.50 | 69.19 | 8.80 | * |
| Σ | 401.20 | 6.83 | 397.28 | 6.13 | * |
| ANSM _e /Nme | 57.18 | 2.27 | 56.04 | 2.33 | * |
| Jarabak | 60.62 | 4.50 | 63.67 | 4.29 | ** |
| Sar/Ramus | 83.75 | 10.23 | 80.52 | 10.40 | |
| PNS-ad1 | 21.95 | 5.83 | 24.83 | 3.91 | * |
| PNS-ad2 | 17.20 | 5.30 | 19.22 | 2.79 | * |
| OAW1 | 9.18 | 2.96 | 10.63 | 2.62 | * |
| OAW2 | 10.58 | 4.56 | 12.15 | 5.57 | |
| OAW3 | 12.52 | 3.44 | 13.85 | 2.76 | |

^a * $P < .05$, ** $P < .01$.

the two adenoideotomy groups were combined for subsequent analysis.

Table 3 displays the comparison results of adenoideotomy and control samples. There was a statistically significant difference in the following parameters: SNGoMe ($P < .01$), PPGoMe ($P < .01$), gonial angle ($P < .05$), gonial ratio ($P < .05$), Σ ($P < .05$), ANSM_e/Nme ($P < .05$), Jarabak ($P < .01$), PNS-ad1 ($P < .05$), PNS-ad2 ($P < .05$), and OAW1 ($P < .05$). Increased skeletal parameters such as SNGoMe, PPGoMe, gonial angle, Σ, ANSM_e/Nme and decreased gonial ratio and Jarabak ratio showed that excessive vertical growth of the adenoideotomy group is evident. Airway measurements indicated that the upper airway of the adenoideotomy group is narrower than that the clear airway group.

DISCUSSION

This study was begun on the hypothesis that the comparison of the craniofacial characteristics of two groups of patients who were operated for adenoids at different stages of their growth and development would give statistically significant differences in the related parameters. Interestingly, however, except in one parameter there were no vertical growth pattern differences between the two groups of children who had adenoidectomy before and after four years of age. This result may point out that up to four years may be a late time in growth to see significant changes after the removal of the excessive lymphoid tissue. There was no statistically significant distinction between the two groups except for ANSMe/Nme parameter, which may be explained by Linder-Aronson's¹² remark that increases in lower anterior face height is independent of other skeletal units and dependent upon the growth direction of the mandible and neuromuscular factors influencing mandibular posture, such as mouth breathing and head posture. The few patients involved in the first group may also be responsible for the absence of a difference between the groups.

When the two adenoidectomy groups were combined and compared with the control group, the adenoidectomy group showed a more vertically directed growth pattern, which is parallel to Linder-Aronson's findings.^{1,11} This clear distinction between groups, ie, the persistent vertical growth model observed in the adenoidectomy group, is probably the outcome of the late date of operation. Because a substantial percentage of facial growth had taken place in most of our patients before the time of operation, there were significant differences between the operated subjects and the controls.

Some authors suggest that adenoids are organs of the immune system and that they should not be removed below the age of eight unless there are absolute indications.²⁹ Likewise, the postoperative management of patients younger than 36 months undergoing adenoidectomy has been the subject of many debates.³⁰ Although the literature questions the safety of ambulatory adenotonsillectomy in children younger than five years, it is a safe procedure with the support of a well-informed and reliable caretaker.³¹ Adenotonsillar hypertrophy, causing upper airway obstruction, may lead to pulmonary alveolar hypoventilation, pulmonary hypertension and cor pulmonale. Early recognition and removal of enlarged lymphoid tissue should be considered for relief of these symptoms.³²⁻³⁵ An adenoidectomy has changed some allergic states positively³⁶ and resolved established (otitis media effusive) OME in 36-46% of cases in children.³⁷ A significant improvement of the mucociliary clearance was shown by Ranga et al³⁸ after adenoidectomy. Therefore, in the long run an earlier operation may improve the patient's quality of life.

We have not included any respiratory measurements using rhinomanometry or nasal respiratory resistances, which represent the cross-sectional area of the tube shaped upper

airway. These measurements reveal anatomic relationships, but they do not indicate a person's breathing pattern. They do not indicate what percentage of the tidal volume is actually respired nasally and orally.²⁵ Therefore, we have used only pharyngeal cephalometric measurements for the evaluation of facial characteristics.

CONCLUSION

This investigation should be considered as a pilot study because of the sample size and its cross-sectional retrospective nature. The longitudinal monitoring of children who had adenoidectomies at very early ages will probably give very precious data towards understanding the mechanisms of the development of long faces.

REFERENCES

1. Linder-Aronson S. Adenoids: their effect on mode of breathing and nasal airflow and their relationship to characteristics of the facial skeleton and the dentition. *Acta Otolaryngol Suppl.* 1970; 265:1-132.
2. Harvold EP, Vargervik K, Chierici G. Primate experiments on oral sensations and dental malocclusions. *Am J Orthod.* 1973;63:494-508.
3. Harvold EP, Tomer BS, Chierici G, Vargervik K. Primate experiments in oral respiration. *Am J Orthod.* 1981;79:359-372.
4. Humphreys HF, Leighton BC. A survey of anteroposterior abnormalities of the jaws in children between the ages of two and five and a half years of age. *Br Dent J.* 1950;88:3-15.
5. Sarver DM. *Quantitative evaluation of nasal airflow in relation to facial morphology* [master's thesis]. Department of Orthodontics, School of Dentistry, University of North Carolina; 1979.
6. Woodside DG, Linder-Aronson S, Lundström A. Mandibular growth direction following adenoidectomy. *Am J Orthod Dentofacial Orthop.* 1986;89:273-284.
7. Hannuksela A. The effect of moderate and severe atopy on the facial skeleton. *Eur J Orthod.* 1981;3:187-193.
8. Watson RM, Warren DW, Fischer ND. Nasal resistance, skeletal classification and mouth breathing in orthodontic patients. *Am J Orthod.* 1968;54:367-379.
9. Rasmus RM, Jacobs RM. Mouth breathing and malocclusion: quantitative technique for measurement of oral and nasal airflow velocities. *Angle Orthod.* 1969;39:296-302.
10. Linder-Aronson S. Effects of adenoidectomy on dentition and nasopharynx. *Am J Orthod.* 1974;65:1-15.
11. Linder-Aronson S. Effects of adenoidectomy on the dentition and facial skeleton over a period of five years. In: Cook JT, ed. *Transactions of the Third International Orthodontic Congress.* St Louis, Mo: The CV Mosby Co; 1975.
12. Woodside DG, Linder-Aronson S. The channelization of upper and lower anterior face heights compared to population standard in males between ages 6 to 20 years. *Eur J Orthod.* 1979;1:25-40.
13. Bresolin D, Shapiro PA, Shapiro GC, Chaptko MK, Dassel S. Mouth breathing in children; its relationship to dentofacial development. *Am J Orthod.* 1983;83:334-340.
14. Wenzel A, Henrikson T, Melsen B. Nasal respiratory resistance and head posture: effect of intranasal corticosteroid (Budesonide) in children with asthma and perennial rhinitis. *Am J Orthod.* 1983; 84:422-426.
15. Behlfelt K, Linder-Aronson S. The effect of enlarged tonsils and tonsillectomy on the dentition and facial skeleton. *Swed Dent J Suppl.* 1990;42.

16. Linder-Aronson S, Bäckström A. A comparison between mouth and nose breathers with respect to occlusion and facial dimensions. *Odontol Revy*. 1960;11:343–376.
17. Tarvonen P, Koski K. Craniofacial skeleton of 7-year-old children with enlarged adenoids. *Am J Orthod Dentofacial Orthop*. 1987;91:300–304.
18. Woodside DG, Linder-Aronson S, Lundström A, McWilliam J. Mandibular and maxillary growth after changed mode of breathing. *Am J Orthod Dentofacial Orthop*. 1991;100:1–18.
19. Gwynne-Evans E, Ballard CF. Discussion on the mouth-breather. *Proc R Soc Med*. 1959;51:279–285.
20. Leech, HL. A clinical analysis of orofacial morphology and behavior of 500 patients attending an upper respiratory research clinic. *Dent Pract*. 1958;9:57–68.
21. Vig PS, Sarver DM, Hall DJ, Warren DW. Quantitative evaluation of nasal airflow in relation to facial morphology. *Am J Orthod*. 1981;79:263–272.
22. Martin R, Vig PS, Warren DW. Nasal resistance and vertical dentofacial features. *J Dent Res*. IADR Abstract, 1981.
23. Schwartzwald R, Rothschild D, Goldsmith D. Vertical facial excess with choanal atresia: Evaluation and treatment considerations. Presented before Committee on Scientific Sessions, American Association of Oral and Maxillofacial Surgeons 62nd annual meeting, 1980.
24. Meredith HV. Growth in head width during the first twelve years of life. *Pediatrics*. 1953;12:411–429.
25. O’Ryan FS, Gallagher DM, LaBanc JP, Epker BN. Relation between nasorespiratory function and dentofacial morphology: a review. *Am J Orthod Dentofacial Orthop*. 1982;403–410.
26. Solow B, Tallgren A. Head posture and craniofacial morphology. *Am J Phys Anthropol*. 1976;44:417–436.
27. Linder-Aronson S, Henrikson CO. Radiocephalometric analysis of anteroposterior nasopharyngeal dimensions in 6 to 12 year old mouth breathers compared with nose breathers. *Orl J (Otorhinolaryngol Rel Spec)*. 1973;35:19–29.
28. Hellsing E. Changes in the pharyngeal airway in relation to extension of the head. *Eur J Orthod*. 1989;11:359–365.
29. Ying MD. Immunological basis of indications for tonsillectomy and adenoidectomy. *Acta Otolaryngol Suppl*. 1988;454:279–285.
30. Shapiro NL, Seid AB, Pransky SM, Kearns DB, Magit AE, Silva P. Adenotonsillectomy in the very young patient: cost analysis of two methods of postoperative care. *Int J Pediatr Otorhinolaryngol*. 1999;48(2):109–115.
31. MacCallum PL, MacRae DL, Sukerman S, MacRae E. Ambulatory adenotonsillectomy in children less than 5 years of age. *J Otolaryngol*. 2001;30(2):75–78.
32. Miman MC, Kirazli T, Ozyurek R. Doppler echocardiography in adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol*. 2000;54(1):21–26.
33. Djalilian M, Kern EB, Brown HA, Facer GW, Stickler GB, Weidman WH, O’Connell EJ. Hypoventilation secondary to chronic upper airway obstruction in childhood. *Mayo Clin Proc*. 1975;50(1):11–14.
34. Karanov J, Minic P, Subarevic V, Baljosevic I. Cor pulmonale caused by hypertrophic adenoid glands and tonsils: indications for tonsillectomy and adenoidectomy in a 2-year-old child. *Srp Arh Celok Lek*. 2000;128(5–6):208–210.
35. Sdralis T, Berkowitz RG. Early adenotonsillectomy for relief of acute upper airway obstruction due to acute tonsillitis in children. *Int J Otolaryngol*. 1996;35(1):25–29.
36. Saito H, Asakura K, Hata M, Kataura A, Morimoto K. Does adenotonsillectomy affect the course of bronchial asthma and nasal allergy. *Acta Otolaryngol Suppl*. 1996;523:212–215.
37. Maw AR. Chronic otitis media with effusion and adeno-tonsillectomy—a prospective randomized controlled study. *Int J Pediatr Otolaryngol*. 1983;6(3):239–246.
38. Ranga RK, Singh J, Gera A, Yadav J. Nasal mucociliary clearance in adenotonsillar hypertrophy. *Indian J Pediatr*. 2000;67(9):651–652.