# **Original Article**

# Dentofacial Characteristics of Chinese Obstructive Sleep Apnea Patients in Relation to Obesity and Severity

# Huie Ming Hou<sup>a</sup>; Urban Hägg<sup>b</sup>; Kim Sam<sup>c</sup>; A.B.M. Rabie<sup>d</sup>; Ricky W.K. Wong<sup>e</sup>; B. Lam<sup>f</sup>; Mary S. Ip<sup>g</sup>

#### ABSTRACT

**Objective:** To evaluate dentofacial characteristics in relation to obesity and degree of severity of obstructive sleep apnea (OSA) in male Chinese patients and to elucidate the relationship between demographic parameters (age, body weight, height, and body mass index [BMI]) cephalometric parameters and OSA in these subjects.

**Materials and Methods:** Lateral cephalograms of 121 Chinese male patients in natural head posture were obtained. Based on BMI value, the patients were divided into three groups. Based on apnea-hypopnea index (AHI) value, the patients were divided into a mild-to-moderate and a severe group.

**Results:** The hyoid position and soft palate length were significantly different among the three obesity groups. Soft palate length was significantly longer (P < .01) in the severe OSA group than in the mild-to-moderate OSA group. Tongue base was significantly more inferiorly placed (P < .05) in the severe OSA group than in the mild-to-moderate OSA group. Craniocervical extension was significantly increased (P < .05) in the severe OSA group. Statistically significant differences were found among the three obesity groups in mandibular length, mandibular body length, maxillary length, anterior cranial base length, and overbite. The multiple stepwise linear regression analysis identified body weight, lower posterior facial height, mandibular body length, craniocervical extension, and sella-hyoid distance as the significant predictive variables for AHI.

**Conclusions:** This study revealed the existence of craniofacial and upper airway soft tissue differences in relation to obesity and severity of OSA among male Chinese OSA patients. Body weight and certain cephalometric parameters were significant predictors of OSA in Chinese male subjects.

KEY WORDS: Obstructive sleep apnea; Cephalometrics; Obesity

#### INTRODUCTION

Obstructive sleep apnea (OSA) is a common disorder, but its pathogenesis is not fully understood. Several cephalometric differences between OSA patients and control samples have been reported. These include mandibular deficiency, bimaxillary retrusion, short cranial base, reduced cranial base angle, mandibular length, increased lower anterior facial height, maxillomandibular plane angle, craniocervical angulation, inferiorly positioned hyoid, and enlarged soft palate.<sup>1–9</sup>

<sup>&</sup>lt;sup>a</sup> Master's student in Orthodontics, Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, People's Republic of China.

<sup>&</sup>lt;sup>b</sup> Chair and Professor in Orthodontics, Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, People's Republic of China.

<sup>°</sup> Part-time Lecturer in Orthodontics, Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, People's Republic of China.

<sup>&</sup>lt;sup>d</sup> Professor in Orthodontics, Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, People's Republic of China.

Accepted: December 2005. Submitted: August 2005. © 2006 by The EH Angle Education and Research Foundation, Inc.

<sup>&</sup>lt;sup>e</sup> Assistant Professor in Orthodontics, Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, People's Republic of China.

<sup>&</sup>lt;sup>f</sup> Medical Doctor, Faculty of Medicine, The University of Hong Kong, Queen Mary Hospital, Hong Kong SAR, People's Republic of China.

<sup>&</sup>lt;sup>9</sup> Chair and Professor in Medicine, Faculty of Medicine, The University of Hong Kong, Queen Mary Hospital, Hong Kong SAR, People's Republic of China.

Corresponding author: Dr. U. Hägg, Orthodontics, Faculty of Dentistry, The University of Hong Kong, Prince Philip Dental Hospital, 34 Hospital Road, Hong Kong SAR, China (e-mail: euohagg@hkusua.hku.hk)

In general, most Asian OSA patients seem to have a lower body mass index (BMI) than their Caucasian counterparts who possess a similar degree of OSA severity.<sup>10–13</sup> There may be differences in obesity and variability in the craniofacial anatomy as risk factors for OSA in these two groups.<sup>13,14</sup> Few English-language cephalometric studies on Chinese OSA subjects have been reported, particularly in relation to obesity and severity of OSA.<sup>15–17</sup> The aims of this study were to evaluate dentofacial characteristics in relation to obesity and degree of severity of OSA in male Chinese patients and to elucidate the relationship between demographic parameters (age, body weight, height, and BMI), cephalometric parameters, and OSA in these subjects.

#### MATERIALS AND METHODS

A total of 121 Chinese male patients with baseline lateral cephalometric radiographs were recruited for this study from a pool of 196 consecutive OSA patients (156 males and 40 females) referred for oral mandibular advancement device (MAD) therapy. Thirty-five of the excluded males were considered not suitable for MAD treatment and no baseline records were obtained. Patients' sleep apnea problems were confirmed by overnight polysomnography and OSA severity was determined by the apnea-hypopnea index (AHI) obtained from the sleep study. Baseline body weight and height were recorded for BMI calculation. Of the 121 patients, 4 patients did not have an AHI value and 16 patients did not have a BMI value. BMI was calculated as follows:

 $BMI = weight (kg) \div height^2 (m)$ 

On the basis of BMI value, the patients were subdivided into three groups: (1) normal-weight (BMI < 25); (2) overweight (25  $\leq$  BMI < 30); and (3) obese (BMI  $\geq$  30). On the basis of AHI value, the patients were subdivided in two groups: (1) mild-to-moderate OSA (AHI < 30); and (2) severe OSA (AHI  $\geq$  30).

# Cephalometric Radiograph and Statistical Analysis

Baseline lateral cephalometric radiographs were obtained with the patients standing upright in natural head posture. The cephalometric landmarks and measurements used in this study are outlined in Figure 1 and Table 1. Computer Assisted Simulation System for Orthognathic Surgery (CASSOS, SoftEnable Technology Limited, Hong Kong SAR China) computer software was used to digitize the lateral cephalometric radiographs, and all radiographs were digitized by the same operator.

The method used for the cephalometric analysis



Figure 1. Cephalometric (a) landmarks, (b) linear measurements, and (c) angular measurements used in this study.

was based on the values obtained from two tracings carried out with an interval of at least 2 weeks. The mean values from the two tracings were used for statistical analyses, which were done using the SPSS 11.0 software package (SPSS, Chicago, ILL). Test for statistical significance was performed with one-way ANOVA for comparison among the normal-weight, Downloaded from https://prime-pdf-watermark.prime-prod.pubfactory.com/ at 2025-05-14 via free access

Table 1. Cephalometric Landmarks and Measurements Used in Figure 1

Variable	Definition
Landmark	
S	Center of the sella turcica.
Ν	Nasion, the deepest point in the concavity of the nasofrontal suture.
Ва	Basion, the most inferior point on the anterior margin of the foramen magnum.
Po	Porion, the most superior point of the bony external auditory meatus.
Or	Orbitale, the most inferior point on the infraorbital margin.
ANS	Anterior nasal spine.
PNS	Posterior nasal spine.
A	A point, the deepest point in the concavity of the anterior maxilla between the ante- rior nasal spine and the alveolar crest.
В	B point, the deepest point in the concavity of the anterior mandible between the alveolar crest and pogonion.
ls	Upper incisor tip.
li	Lower incisor tip.
Ms	Mesial buccal cusp tip of upper first molar.
Cd	Condylion, the most posterosuperior point of the condylar head.
Go	Gonion, the most posterioinferior point on the angle of the mandible.
Gn	Gnathion, the most anteroinferior point on the bony chin.
Me	Menton, the most inferior point on the body chin.
Rgn	Retrognathia, the most posterior point of symphysis.
U	The tip of the uvula.
V	Vallecula, the most posteroinferior base of the tongue.
H	The most superior and anterior point on the body of the hyoid bone.
c2sp	The most superior posterior point of the second cervical vertebra.
c2ip	The most inferior posterior point of the second cervical vertebra.
c4la	The most inferior anterior point of the fourth cervical vertebra.
	The most interior posterior point of the fourth cervical vertebra.
	Frankfort nonzontal plane, the line joining Po and Or.
	Manulpular plane, the line joining DNS and ANS
Mixel	The line ioning is and Me (take the second moler if the first moler is missing and
	take the premolar if both molars are missing).
OPT	Odomola process langent, the line joining c2sp and c2p.
Hyoid position	
Rgn-H (mm)	The distance between Rgn and H.
H-MnPI (mm)	The perpendicular distance from H to the MnPI.
H-c4ia (mm)	The distance between H and c4ia.
S-H (mm)	The distance between S and H.
Tongue base position	
PNS-V (mm)	Length of pharynx, the distance between PNS and V.
V-FH (mm)	Vertical position of V, the perpendicular distance between V and Frankfort horizon- tal plane.
V-c4ia (mm)	Horizontal position of V, the distance between V to c4ia, parallel to Frankfort hori- zontal plane.
Soft palate	
PNS-U (mm)	Soft palate length, the distance between PNS and U.
Soft palate angle (°)	The angle between soft palate length (U-PNS) and MxPl.
Craniocervical extension	
OPT-SN (°)	The angle between the OPT and the sella-nasion (S-N) line
c2sp-c4ip-SN (°)	The angle between the c2sp-c4ip line and the S-N line.

overweight, and obese groups. Student's *t*-test was used for comparison between the mild-to-moderate and severe OSA groups. *P* values less than .05 were considered significant. Linear regression analysis with a forward selection stepwise procedure was used to evaluate the association between AHI (dependent factor) and age, body weight, height, BMI, and cephalometric variables in Figure 1 (independent factor).

#### Method Error

The method error study was carried out by repeating the digitization process for ten randomly-selected ra-

Table 1. C	ontinued.
------------	-----------

Variable	Definition				
Craniofacial structures					
NSBa (°)	Skull base angle, the angle between the nasion-sella (N-S) line and the sella-basi- on (S-Ba) line.				
SNA (°)	The angle between the S-N line and the Nasion-A (N-A) line.				
SNB (°)	The angle between the S-N line and the Nasion-B (N-B) line.				
ANB (°)	The angle between the N-A line and the N-B line.				
MnPl/SN (°)	Mandibular plane angle, the angle between the MnPI and the S-N line.				
MxPI/SN (°)	Maxillary plane angle, the angle between the MxPI and the S-N line.				
Overjet (mm)	The distance between the Is and Ii, parallel to the upper occlusal plane (positive if upper incisor is in front of the lower incisor, negative if lower incisor is in front of the upper incisor).				
Overbite (mm)	The distance between Is and Ii, perpendicular to the upper occlusal plane (positive if there is overlapping, negative if there is open bite).				
Upper posterior facial height (UPFH) (mm)	The distance from S to MxPI, along the S-Go line.				
Lower posterior facial height (LPFH) (mm)	The distance from MxPI to Go, along the S-Go line.				
Total posterior facial height (TPFH) (mm)	The distance from S to Go.				
Upper anterior facial height (UAFH) (mm)	The distance from N to MxPI, along the N-Me line.				
Lower anterior facial height (LAFH) (mm)	The distance from MxPI to Me, along the N-Me line.				
Total anterior facial height (TAFH) (mm)	The distance from N to Me.				
Mandibular length (mm)	The distance between Cd and Gn.				
Ramus length (mm)	The distance between Cd and Go.				
Body length (mm)	The distance between Go and Me.				
Maxillary length (mm)	The distance between ANS and PNS.				
Anterior cranial base length (SN) (mm)	The distance between S and N.				

Table 2. Demographic Data of Normal-Weight (n = 38; 36%), Overweight (n = 43; 41%), and Obese (n = 24; 23%) Male OSA Patients

	Normal-Weight			Overweight			Obese				
Variable	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range	Р	
Age (y)	48.7	10.3	20.5-66.6	47.1	9.2	27.8–71.4	42.6	7.7	27.7-62.1	.047*	
Body weight (kg)	67.8	5.8	59.1-86.5	77.4	5.8	65.5-92.0	104.5	25.3	80.5-197.0	<.001***	
Height (cm)	171	0.1	159–195	170	0.1	156–178	171	0.1	163–185	.663	
BMI (kg/m <sup>2</sup> )	23.2	1.7	16.0-24.9	26.9	1.4	25.0-29.9	35.7	8.1	30.1-67.4	<.001***	
AHI	24.4	16.0	4.7–70.1	28.5	18.5	3.3–65.3	45.8	24.2	9.4-88.0	<.001***	

<sup>a</sup> OSA indicates obstructive sleep apnea; BMI, body mass index, and AHI, apnea-hypopnea index; \* P < .05; \*\*\* P < .001.

diographs, again at an interval of at least 2 weeks. Mean values from the first and second tracings together with mean values from the third and fourth tracings were used for the method error determination. The method error was determined with the formula

$$\mathsf{SE} = \sqrt{\sum d^2/2n}$$

where  $\Sigma d^2$  is the sum of the squared differences between the two mean values, and *n* is the number of double measurements.

For linear measurements the method error was determined as 0.4 mm (P > .1) and for angular measurements 0.5° (P > .1). Both of these were statistically insignificant. It can therefore be concluded that no systematic error occurred for landmark identification or cephalometric measurements.

Approval for the study was obtained from the ethics committee of the Faculty of Dentistry, University of Hong Kong.

#### RESULTS

There were significant differences in age, weight, BMI, and AHI among the three obesity groups (Table 2). The obese group was the youngest and had the highest AHI value. Between the mild-to-moderate and severe OSA groups, there were significant differences in weight, BMI, and AHI (Table 3). The severe OSA group was younger and more obese than the mild-tomoderate OSA group.

The hyoid bone position (Figure 1b) as measured from Rgn, MnPl, c4ia, and S was significantly different among the three obesity groups (Table 4). The hyoid bone was most inferiorly placed in the obese group, whereas it was most superiorly placed in the normalweight group. There was no statistically significant difference in hyoid position between the mild-to-moderate and severe OSA groups (Table 5). As for tongue base position, statistically significant differences were found only in the PNS-V measurement between the Downloaded from https://prime-pdf-watermark.prime-prod.pubfactory.com/ at 2025-05-14 via free access

		Mild-to-Mod	erate				
Variable	Mean	SD	Range	Mean	SD	Range	Р
Age (y)	46.1	11.5	20.5-79.2	47.2	8.5	27.8-65.2	.558
Body weight (kg)	74.8	10.8	61.5-113.0	86.6	24.5	59.1-197.0	.003**
Height (cm)	170	0.1	158–181	171	0.1	156–195	.528
BMI (kg/m <sup>2</sup> )	25.9	3.6	19.4-40.5	29.7	7.9	16.0-67.4	.004**
AHI	15.7	7.6	3.3–29.9	49.2	16.1	30.0-88.0	<.001***

Table 3. Demographic Data of Mild-to-Moderate (n = 64; 55%) and Severe (n = 53; 45%) Male OSA Patients

<sup>a</sup> OSA indicates obstructive sleep apnea; BMI, body mass index, and AHI, apnea-hypopnea index; \*\* *P* < .01; \*\*\* *P* < .001.

Table 4.	Cephalometric	Comparison o	f Normal-Weight (n =	- 38),	Overweight (n = $43$ ),	and Obese	(n = 24)	Male OSA Patients
----------	---------------	--------------	----------------------	--------	-------------------------	-----------	----------	-------------------

	Normal-Weight		Overweight		Obese		
Variable	Mean	SD	Mean	SD	Mean	SD	Р
Hyoid position							
Rgn-H (mm)	42.4	4.6	43.9	6.0	48.0	4.9	<.001***
H-MnPl (mm)	22.6	4.9	23.5	5.4	26.4	5.5	.014
H-c4ia (mm)	45.2	4.2	45.9	4.9	49.8	5.7	.011
S-H (mm)	128.6	7.8	130.7	6.7	132.4	9.6	.167
Tongue base position							
PNS-V (mm)	84.1	6.2	85.3	6.4	87.7	7.6	.063
V-FH (mm)	107.3	6.7	109.4	6.4	110.4	8.9	.165
V-c4ia (mm)	107.3	6.7	109.4	6.4	110.4	8.9	.666
Soft palate							
PNS-U (mm)	47.0	4.3	47.5	3.7	50.2	4.2	.004
Soft palate angle (°)	125.9	4.8	126.1	5.6	124.7	5.7	.526
Craniocervical extension							
OPT-SN (°)	110.7	8.1	109.8	7.2	113.5	6.4	.093
c2sp.c4ip-SN (°)	113.3	7.1	112.6	6.2	115.7	6.5	.121
Craniofacial structures							
NSBA (°)	128.3	5.5	127.3	5.7	130.3	6.5	.139
SNA (°)	81.0	4.3	82.4	4.2	82.3	4.0	.391
SNB (°)	77.2	4.0	78.4	3.8	79.3	4.4	.221
ANB (°)	4.0	2.8	4.0	3.0	3.0	2.2	.353
MnPl/SN (°)	36.6	6.5	34.3	7.3	35.4	5.6	.449
MxPI/SN (°)	10.6	3.1	9.5	3.9	10.8	3.0	.262
Overjet (mm)	4.6	2.7	4.7	2.7	3.1	2.4	.046
Overbite (mm)	3.4	2.2	2.7	1.7	2.1	1.8	.020
UPFH (mm)	48.2	3.6	49.3	3.9	47.7	3.7	.207
LPFH (mm)	43.0	3.9	43.9	4.7	44.8	5.5	.309
TPFH (mm)	91.2	5.4	93.2	5.2	92.5	6.2	.318
UAFH (mm)	61.8	3.0	61.5	3.3	61.8	2.9	.917
LAFH (mm)	75.1	5.4	75.2	5.9	76.8	4.4	.357
TAFH (mm)	136.8	6.9	136.8	7.8	138.6	6.0	.433
Mandibular length (mm)	124.7	5.6	126.8	6.3	130.8	6.5	.001
Ramus length (mm)	66.5	4.2	68.2	4.4	68.6	4.6	.165
Mandibular body length (mm)	72.8	5.1	74.9	5.5	76.8	4.8	.012
Maxillary length (mm)	50.9	2.7	53.0	2.9	54.2	3.2	<.001***
SN length (mm)	73.2	2.7	74.2	3.3	75.3	2.5	.028

° OSA indicates obstructive sleep apnea; \* P < .05; \*\* P < .01; \*\*\* P < .001.

mild-to-moderate OSA and severe OSA groups, where the tongue base was 3.0 mm more inferiorly placed in the severe OSA group (P < .05). No statistically significant difference in the tongue base position was found among the three obesity groups. For soft palate measurements, soft palate length (PNS-U) was significantly different for the three obesity groups, with the longest measurement in the obese group. It was also significantly longer in the severe OSA group than the mild-to-moderate OSA group (P < .01).

Table 5.	Cephalometric Con	nparison of Mild-to-Mod	erate ( $n = 64$ )	and Severe (	n = 53	) Male OSA	Patients
----------	-------------------	-------------------------	--------------------	--------------	--------	------------	----------

	Mild-to-M	oderate	Seve		
Variable	Mean	SD	Mean	SD	P
Hyoid position					
Rgn-H (mm)	43.7	5.0	44.4	6.3	.508
H-MdPI (mm)	23.2	5.0	24.6	5.7	.156
H-c4ia (mm)	45.8	4.7	47.0	5.5	.257
S-H (mm)	129.4	7.5	131.9	8.3	.086
Tongue base position					
PNS-V (mm)	83.9	5.9	86.9	6.7	.014*
V-FH (mm)	107.6	7.2	110.2	7.0	.057
V-c4ia (mm)	28.1	5.8	28.1	5.6	.957
Soft palate					
PNS-U (mm)	47.0	4.2	49.0	3.7	.009**
Soft palate angle (°)	126.8	4.6	125.0	6.0	.081
Craniocervical extension					
OPT-SN (°)	109.6	6.6	112.5	7.9	.032*
c2sp.c4ip-SN (°)	112.3	5.9	114.9	7.0	.036*
Craniofacial structures					
NSBA (°)	128.7	6.0	128.1	5.8	.606
SNA (°)	81.7	4.1	82.0	4.3	.765
SNB (°)	78.2	3.8	77.9	4.6	.708
ANB (°)	3.6	2.7	4.0	2.6	.410
MnPI/SN (°)	35.1	5.8	36.0	7.6	.529
MxPI/SN (°)	9.7	3.2	10.6	3.5	.151
Overjet (mm)	4.3	3.0	4.1	2.3	.693
Overbite (mm)	2.7	2.2	2.7	1.8	.913
UPFH (mm)	49.0	3.3	48.3	4.3	.287
LPFH (mm)	42.9	3.7	44.7	5.3	.035*
TPFH (mm)	92.0	4.7	93.0	6.1	.305
UAFH (mm)	61.6	2.8	61.9	3.3	.527
LAFH (mm)	75.2	5.0	76.3	6.0	.309
TAFH (mm)	136.7	6.4	138.2	7.9	.278
Mandibular length (mm)	126.9	5.8	127.0	7.2	.919
Ramus length (mm)	67.3	4.2	68.5	4.5	.141
Mandibular body length (mm)	74.8	5.1	73.9	5.6	.398
Maxillary length (mm)	52.3	3.1	52.8	3.3	.440
SN length (mm)	74.2	2.7	74.1	3.2	.830

<sup>a</sup> OSA indicates obstructive sleep apnea; \* P < .05; \*\* P < .01.

Craniocervical extension measurements (OPT/SN and c2sp.c4ip/SN angles) (Figure 1c) differed significantly between the mild-to-moderate and severe OSA groups (Table 5), but not among the three obesity groups (Table 4). The mean OPT/SN angle in severe OSA group was 3°, (P < .05) higher than that found in the mild-to-moderate OSA group. The mean c2sp-c4ip-SN angle in the severe OSA group was also 3°, higher than that found in the mild-to-moderate OSA group (P < .05).

The only significant difference in the measurements of the craniofacial structures between the mild-to-moderate and severe OSA groups was in lower posterior facial height (LPFH) (Table 5) where the mean was 1.8 mm (P < .05) longer in the severe OSA group. There were statistically significant differences in mandibular length (Cd-Gn), mandibular body length (GoMe), maxillary length (ANS-PNS), sella-nasion length (S-N), overjet, and overbite between the three obesity groups. The obese group had the greatest Cd-Gn, Go-Me, ANS-PNS, and S-N, whereas the normal-weight group had the shortest lengths for these measurements. For overjet and overbite, the normal-weight group had the most protrusive teeth and deepest bite among the three groups.

For all the OSA patients, the multiple stepwise linear regression model for AHI ( $R^2 = 0.434$ , P < .001) identified the body weight (partial  $R^2 = 0.162$ , P < .001), mandibular body length (partial  $R^2 = 0.114$ , P < .01), LPFH (partial  $R^2 = 0.076$ , P < 0.01), c2sp-c4ip-SN angle (partial  $R^2 = 0.044$ , P < .01) and S-H (partial  $R^2 = 0.038$ , P < .05) as the significant predictive variables (Table 6).

**Table 6.** Linear Regression Model with AHI as Dependent Factorand Age, Weight, Height, BMI, and Cephalometric Variables (Figure1) as Independent Factors

<i>,</i> ,			
Variable	Coefficient	Standard error	Р
Body weight (kg)	0.383	0.092	.001
Body length (mm)	-1.192	0.346	.001
LPFH (mm)	1.399	0.410	.001
c2sp.c4ip-SN (°)	0.761	0.280	.008
S-H (mm)	0.557	0.249	.028
Constant	-132.287	54.196	.017

<sup>a</sup> AHI indicates apnea-hypopnea index; BMI, body mass index.

# DISCUSSION

The demographic data from this study (Table 2) show that the obese group was more severely affected by OSA (increased AHI). Previous studies have also found an association between obesity and OSA.<sup>11,18</sup> The BMI grouping (Table 2) and AHI grouping (Table 3) show statistically significant differences in BMI and weight, but not in height, indicating that weight is a more important factor in determining the statistical significance of BMI. Furthermore, in the multiple stepwise linear regression analysis, body weight was found as the dominant significant predictor of AHI.

This study noted that OSA tended to occur at a younger age in the obese group, but previous studies have indicated that the prevalence of OSA seems to increase in midlife, with higher prevalence among older age groups.<sup>19,20</sup> The risk of OSA is much higher with increased obesity, and this probably explains why an earlier occurrence of OSA was observed in the obese group in this study (Table 2). Because obesity involving all age groups is becoming an increasingly serious problem in many affluent societies, particularly the United States,<sup>21</sup> it is likely that a growing number of young obese people will be affected by OSA in these societies in the years to come.

This study found that normal-weight OSA patients had significantly greater anteroposterior facial compression than obese OSA patients (Table 4). This manifested as a shorter mandible (mandibular length and body length), shorter maxilla (maxillary length), and shorter anterior cranial base length (SN length). These cephalometric findings are consistent with the results of some earlier studies.<sup>17,22–24</sup>

The multiple stepwise linear regression analysis revealed that the mandibular body length was a significant predictor for AHI in this study (Table 6). In another study on Chinese subjects, it was found that retroposition of the mandible was associated with OSA, particularly with more severe OSA.<sup>25</sup> Some studies have also reported that normal-weight OSA patients have more craniofacial structure abnormalities than their obese counterparts, whereas obese OSA patients tend to have more normal craniofacial structures but more upper airway soft tissue abnormalities.<sup>22-24,26</sup>

The hyoid bone is important as anchorage for the tongue muscles, and its position partly affects the shape, size and position of the tongue. Previous studies have shown that the hyoid bone is more inferiorly positioned in OSA patients than in non-OSA subjects.<sup>1,2,6</sup> However, other studies found that the difference in the position of the hyoid bone in nonobese and obese OSA patients was not significant.22,24 By contrast, this study revealed that the hyoid bone was more inferiorly positioned in obese OSA patients. One explanation for this finding might be that the BMI of the obese OSA group in this study (Table 2) was higher than that found in the two earlier studies.22,24 This might have resulted in increased fat deposition in the upper airway soft tissues, forcing the hyoid bone into a more inferior position. The inferior position of the hyoid bone in turn gave the tongue a more upright position, and more tongue mass occupied the hypopharyngeal area.<sup>8</sup> Multiple stepwise linear regression analysis of this study found that the position of hyoid bone from sella was a significant predictor of AHI (Table 6). Similarly, a recent study on Chinese subjects also reported the inferior hyoid position as one of the cephalometric parameters associated with OSA.25

The increased soft palate length of the obese OSA patients in this study could be related to fat deposits caused by increased obesity. Previous studies have also shown upper airway soft-tissue abnormalities such as increased soft palate length in obese OSA patients.<sup>22,24,26,27</sup> The results of this study suggest that the increased soft palate length and inferiorly positioned hyoid bone in obese OSA patients could make them more prone to airway obstruction.

As for the mild-to-moderate OSA group vs the severe OSA group, significant differences were found in soft palate length and tongue base position, but not in hyoid bone position. The severe OSA group had a longer soft palate length and the tongue base was in a more inferior position. These features in the severe OSA group indicate that more soft palate and tongue masses would occupy the upper airway space, and this in turn could make the upper airway more prone to obstruction and might partly account for the increased OSA severity in this sample.

The craniocervical extension has also been related to the severity of OSA.<sup>6,7,28</sup> The craniocervical extension measurements made in this study (OPT/SN and c2sp.c4ip/SN) showed that the head posture differed significantly between the mild-to-moderate OSA group and the severe OSA group. Severe OSA patients had a more extended craniocervical angulation than did mild-to-moderate OSA patients, similar to findings noted in other studies.<sup>6,7,28</sup> The increased craniocervical angle in the OSA patients was a physiological adaptation aimed at maintaining airway adequacy.<sup>7</sup> This probably represents a compensatory physiological adaptation to lift away the base of the tongue and the soft palate from the posterior pharyngeal wall, thereby alleviating the obstruction and maintaining airway adequacy.<sup>7</sup> In addition, in this study multiple stepwise linear regression analysis confirmed that the c2sp.c4ip/SN measurement is one of the significant predictor for AHI.

### CONCLUSIONS

- Obesity and craniofacial and soft tissue abnormalities may aggravate the severity of OSA syndrome in Chinese male subjects.
- Body weight and certain cephalometric parameters are significant predictors of OSA in Chinese male subjects.

## ACKNOWLEDGMENTS

Sincere thanks to Mr Shadow Yeung for his kind assistance with the statistical analyses and Ms Zinnia Pang for her assistance with the manuscript.

## REFERENCES

- Jamieson A, Guilleminault C, Partinen M, Quera-Salva MA. Obstructive sleep apneic patients have craniomandibular abnormalities. *Sleep.* 1986;9:469–477.
- 2. Tsuchiya M, Lowe AA, Pae EK, Fleetham JA. Obstructive sleep apnea subtypes by cluster analysis. *Am J Orthod Dentofacial Orthop.* 1992;101:533–542.
- 3. Lowe AA, Santamaria JD, Fleetham JA, Price C. Facial morphology and obstructive sleep apnea. *Am J Orthod Dentofacial Orthop.* 1986;90:484–491.
- 4. Andersson L, Brattstrom V. Cephalometric analysis of permanently snoring patients with and without obstructive sleep apnea syndrome. *Int J Oral Maxillofac Surg.* 1991;20:159– 162.
- Rivlin J, Hoffstein V, Kalbfleisch J, McNicholas W, Zamel N, Bryan AC. Upper airway morphology in patients with idiopathic obstructive sleep apnea. *Am Rev Respir Dis.* 1984; 129:355–360.
- Tangugsorn V, Skatvedt O, Krogstad O, Lyberg T. Obstructive sleep apnea: a cephalometric study. Part I. Cervicocraniofacial skeletal morphology. *Eur J Orthod.* 1995;17:45– 56.
- Solow B, Ovesen J, Nielsen PW, Wildschiodtz G, Tallgren A. Head posture in obstructive sleep apnea. *Eur J Orthod.* 1993;15:107–114.
- Lyberg T, Krogstad O, Djupesland G. Cephalometric analysis in patients with obstructive sleep apnea syndrome: II. Soft tissue morphology. *J Laryngol Otol.* 1989b;103:293– 297.
- Tangugsorn V, Skatvedt O, Krogstad O, Lyberg T. Obstructive sleep apnea: a cephalometric study. Part II. Uvulo-glossopharyngeal morphology. *Eur J Orthod.* 1995;17:57–67.
- Ip MS, Lam B, Lauder IJ, Tsang KW, Chung KF, Mok YW, Lam WK. A community study of sleep-disordered breathing in middle-aged Chinese men in Hong Kong. *Chest.* 2001; 119:62–69.

- Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med.* 1993;328:1230–1235.
- Bearpark H, Elliott L, Grunstein R, Cullen S, Schneider H, Althaus W, Sullivan C. Snoring and sleep apnea. A population study in Australian men. *Am J Respir Crit Care Med.* 1995;151:1459–1465.
- Li KK, Powell NB, Kushida C, Riley RW, Adornato B, Guilleminault C. A comparison of Asian and white patients with obstructive sleep apnea syndrome. *Laryngoscope.* 1999; 109:1937–1940.
- 14. Liu Y, Lowe AA, Zeng X, Fu M, Fleetham JA. Cephalometric comparisons between Chinese and Caucasian patients with obstructive sleep apnea. *Am J Orthod Dentofacial Orthop.* 2000;117:479–485.
- 15. Ang PK, Sandham A, Tan WC. Craniofacial morphology and head posture in Chinese subjects with obstructive sleep apnea. *Semin Orthod.* 2004;10:90–96.
- Hui DS, Ko FW, Chu AS, et al. Cephalometric assessment of craniofacial morphology in Chinese patients with obstructive sleep apnea. *Respir Med.* 2003;97:640–646.
- Liao YF, Chuang ML, Huang CS, Tsai YY. Upper airway and its surrounding structures in obese and nonobese patients with sleep-disordered breathing. *Laryngoscope*. 2004; 114:1052–1059.
- Davies RJ, Ali NJ, Stradling JR. Neck circumference and other clinical features in the diagnosis of the obstructive sleep apnea syndrome. *Thorax.* 1992;47:101–105.
- Bixler EO, Vgontzas AN, Lin HM, Ten Have T, Rein J, Vela-Bueno A, Kales A. Prevalence of sleep-disordered breathing in women: effects of gender. *Am J Respir Crit Care Med.* 2001;163:608–613.
- Young T, Shahar E, Nieto FJ, et al. Predictors of sleepdisordered breathing in community-dwelling adults: the Sleep Heart Health Study. Arch Intern Med. 2002;162:893– 900.
- Mokdad AH, Serdula MK, Dietz WH, Bowman BA, Marks JS, Koplan JP. The spread of the obesity epidemic in the United States, 1991–1998. *JAMA*. 1999;282:1519–1522.
- Sakakibara H, Tong M, Matsushita K, Hirata M, Konishi Y, Suetsugu S. Cephalometric abnormalities in non-obese and obese patients with obstructive sleep apnea. *Eur Respir J.* 1999;13:403–410.
- Paoli JR, Lauwers F, Lacassagne L, Tiberge M, Dodart L, Boutault F. Craniofacial differences according to the body mass index of patients with obstructive sleep apnea syndrome: cephalometric study in 85 patients. *Br J Oral Maxillofac Surg.* 2001;39:40–45.
- 24. Yu X, Fujimoto K, Urushibata K, Matsuzawa Y, Kubo K. Cephalometric analysis in obese and nonobese patients with obstructive sleep apnea syndrome. *Chest.* 2003;124: 212–218.
- 25. Lam B, Ooi CG, Peh WC, Lauder I, Tsang KW, Lam WK, Ip MS. Computed tomographic evaluation of the role of craniofacial and upper airway morphology in obstructive sleep apnea in Chinese. *Respir Med.* 2004;98:301–307.
- Ferguson KA, Ono T, Lowe AA, Ryan CF, Fleetham JA. The relationship between obesity and craniofacial structure in obstructive sleep apnea. *Chest.* 1995;108:375–381.
- Ito D, Akashiba T, Yamamoto H, Kosaka N, Horie T. Craniofacial abnormalities in Japanese patients with severe obstructive sleep apnea syndrome. *Respirology*. 2001;6:157–161.
- Solow B, Skov S, Ovesen J, Norup PW, Wildschiodtz G. Airway dimensions and head posture in obstructive sleep apnea. *Eur J Orthod.* 1996;18:571–579.