# **Original Article**

# Patients' self-perception of dentofacial attractiveness before and after exposure to facial photographs

## Giulio Alessandri Bonettia; Andrea Albertib; Claudio Sartinic; Serena Incerti Parentib

#### ABSTRACT

**Objective:** To determine whether exposure to pretreatment photographs of themselves influenced patients' self-perception of dentofacial attractiveness and willingness to undergo treatment.

**Materials and Methods:** A total of 100 subjects of white ancestry aged 18 years or older (Study Group) and 100 sex- and age-matched controls (Control Group) were selected. Photographs of the frontal view of the face and the profile view of the face, both at rest and while smiling, were taken of each participant. Only Study Group subjects were given a printed copy of their own images to be examined at home between the initial observation (T0) and a set date an average of 30 days later (T1). Each of the 200 subjects in the study completed a questionnaire at T0 and T1 exploring happiness regarding their smiles and their facial profiles, as well as willingness to undergo treatment. **Results:** At T1 in the Study Group, 50% of subjects responded with a lower opinion of their facial profiles than at T0; 11% and 45% were willing to undergo more comprehensive procedures to change the appearance of their smiles and profiles, respectively. No statistically significant change was seen in questionnaire answers between T0 and T1 for the Control Group.

**Conclusion:** Laypeople generally are not aware of their facial profiles. Pretreatment photograph exposure can increase profile self-awareness, an important factor in reducing the discrepancy between orthodontists' and patients' visual emphasis on dentofacial esthetics. (*Angle Orthod.* 2011;81:517–524.)

KEY WORDS: Profile; Self-perception; Facial photographs; Dentofacial attractiveness

# INTRODUCTION

Modern society has placed a strong emphasis on physical appearance. The desire to improve dentofacial esthetics has been found to be the primary motivation for patients seeking orthodontic care, regardless of structural or functional considerations.<sup>1-4</sup>

Therefore, self-perception of their own dentofacial attractiveness is a key motivational factor for patients seeking orthodontic evaluation and is an important factor in their expectation of treatment outcome.

Corresponding author: Dr Giulio Alessandri Bonetti, Department of Orthodontics, University of Bologna, via San Vitale, 59 Bologna, 40125 Italy

(e-mail: giulio.alessandri@unibo.it)

Accepted: November 2010. Submitted: October 2010.

Published Online: February 7, 2011

However, this self-perception essentially is based on how individuals see themselves in the mirror, with frontal views of the face and smile typically representing their primary concerns.<sup>1,5</sup>

Although orthodontic diagnosis is carried out in three dimensions (transverse, anteroposterior, and vertical), orthodontists place major treatment planning emphasis on the esthetics of the face in profile.<sup>6,7</sup> Because most people cannot characterize their own profile,<sup>1</sup> a difference does exist between orthodontic professionals and the public regarding perceptions of facial profile esthetics.<sup>3,8–17</sup>

The aim of the current study, using a sample of subjects seeking or referred for orthodontic evaluation, was to determine whether exposure to pretreatment smile and profile photographs influenced individuals' self-perception of dentofacial attractiveness and willingness to undergo treatment.

#### MATERIALS AND METHODS

This questionnaire-based study included two groups of subjects: 100 subjects (54 females and 46 males; referred to as the Study Group) who were selected

<sup>&</sup>lt;sup>a</sup> Assistant Professor, Department of Orthodontics, University of Bologna, Bologna, Italy.

<sup>&</sup>lt;sup>b</sup> Resident, Department of Orthodontics, University of Bologna, Bologna, Italy.

<sup>&</sup>lt;sup>o</sup> Biostatistician (MSC), Environmental Protection Agency of Emilia Romagna Region, Epidemiology Unit, Modena, Italy.

 $<sup>{\</sup>scriptstyle \circledcirc}$  2011 by The EH Angle Education and Research Foundation, Inc.

Table 1.	Sample	Distribution	bv	Aae
	oumpic	Distribution	ъy	/ igo

Age, years	Study Group, N° of Subjects	Control Group, N° of Subjects	Total, N° of Subjects
18–24	18	18	36
25–29	10	10	20
30–34	20	20	40
35–39	35	35	70
40–45	11	11	22
46–50	6	6	12
Total, N° of			
subjects	100	100	200

among individuals referred to the Department of Orthodontics within the School of Dentistry, University of Bologna, and 100 additional age- and sex-matched subjects (the Control Group). The study took place between 2005 and 2008. Given the difficulty involved in finding exact age matches, selection of controls was based on a range of  $\pm 6$  months independent of the year of birth. The distribution of subjects' ages is presented in Table 1.

The study protocol, the questionnaire, and the informed consent form were approved by the Institutional Review Board. All subjects agreed to participate in the study and signed a written informed consent, although no explanation of the goals was provided. Subjects received no compensation for their participation.

All participants met the following inclusion criteria: (1) white origin, (2) older than 18 years of age, and (3) seeking or referred for orthodontic evaluation. Exclusion criteria were as follows: (1) previous orthodontic treatment or plastic or orthognathic surgery, and (2) any systemic medical condition that might affect subjects' physical or emotional growth, including psychiatric conditions.

# **Data Collection**

Subjects' responses regarding their perceived dentofacial attractiveness and willingness to undergo treatment were recorded using a specially designed questionnaire. The first section of the questionnaire focused on the demographic data of participants, such as age, gender, marital status, income level, and level of education. Subjects were asked whether they liked the appearance of their smiles (Question #1) and the appearance of their faces in profile (Question #2), using a 4-point scale with 1 as "very much," 2 as "somewhat," 3 as "a little bit," and 4 as "not at all." Finally, subjects were asked which treatment options they would seek to change the appearance of their smiles (Question #3) and their facial profiles (Question #4), using a 4-point scale with 1 as "None," 2 as "Removable orthodontic appliance," 3 as "Fixed orthodontic appliance," and 4 as "Surgery."

After verbal instructions were given, each individual was seated in a quiet area apart from the other subjects. Each was given 10 minutes to anonymously complete the questionnaires, which were marked by numeric codes. Participants were advised that the information collected was confidential and would not affect their future care. They were also given the freedom to leave the study at any time. Two investigators were available to explain the questions and to check the questionnaires for completeness.

After completion of the questionnaire, a dental evaluation was performed by one experienced orthodontist, who conducted all of the clinical examinations. Angle Class was rated on a 4-point scale with 1 as "Class I," 2 as "Class II, Division 1," 3 as "Class II, division II," and 4 as "Class III." Overjet was scored on a 4-point scale with 1 as "Reduced (<2 mm)," 2 as "Normal (2 to 3 mm)," 3 as "Increased (>3 to 6 mm)," and 4 as "Very increased (>6 mm)." Overjet and Angle Class characteristics are reported in Table 2.

Photographs of frontal and profile views of the face, both at rest and while smiling, were taken of each participant using a Nikon Digital SLR Camera D90 (Nikon Photo Products, Tokyo, Japan) under standard conditions. Several photos of each subject were produced, so that natural and unforced neutral facial expressions and smiles could be chosen and subsequently printed.

All subjects filled in the questionnaire twice, separated by an average interval of 30 days (T0, first completion; T1, second completion), and both questionnaires for a particular subject were identified by the same numeric code. During the period between T0 and T1, only subjects in the Study Group were given a printed copy of the photographs of themselves (Figure 1). They were instructed to show the photographs to relatives and friends for discussion.

Table 2. Sample Distribution by Overjet and Angle Class

	Overjet			Angle Class				
	Reduced, <2 mm	Normal, 2–3 mm	Increased, >3–6 mm	Very increased, >6 mm	Class I	Class II, Division 1	Class II, Division 2	Class III
Study Group	3	30	51	16	32	62	3	3
Control Group	4	29	44	23	32	58	6	4
Total	7	59	95	39	64	120	9	7



Figure 1. Representative photographs used in this study: (1) frontal full-face photograph at rest and (2) while smiling; (3) profile; (4) profile at rest and (5) while smiling; and (6) frontal smile. (1) through (3) are full-face photographs, and photographs in (4) through (6) feature particular portions of the face.

#### **Reliability of the Questionnaire**

As a preliminary test, 10 subjects were selected, on a consecutive basis, among individuals referred to the Department of Orthodontics, in full accordance with inclusion and exclusion criteria of the study. To determine the reliability of the questionnaire, subjects were asked to complete it twice, separated by a 2-week interval, with the researcher present and after written consent had been obtained. Although both tests were administered anonymously, correlation of the second questionnaire with the first was possible because of the numeric code. Responses from both questionnaires were coded, and the Kappa statistic (Cohen's Kappa,  $\kappa$ ) was used to assess reliability.

The time taken by each participant to complete the questionnaire at the first completion was recorded by the researcher using a stopwatch to determine the ease of administration. The readability of the questionnaire was assessed using the Flesch Reading Ease Score, the Flesch-Kincaid Grade Level readability statistics, and the Passive Sentences Score formula (Microsoft Word 2007, Microsoft Corporation, Red-mond, Wash).

#### **Statistical Analysis**

As a preliminary analysis, the equivalence between Study and Control Groups was investigated with respect to all examined variables (level of income, marital status, educational level, overjet, and Angle Class). The Kruskal-Wallis nonparametric equality-ofpopulations rank test was used to compare the two groups.

Responses from the first and second completions of the questionnaires were coded. The differences between T0 and T1 were evaluated for each response and compared between groups using a chi-square test.

A logistic regression analysis was applied to adjust for all potential confounders (i.e., age, marital status, level of income, level of education, overjet, and Angle Class). The outcome was a negative change in subjects' opinions between T0 and T1. Statistical analysis was performed using STATA 10 (StataCorp, College Station, Tex).

# RESULTS

#### **Reliability of the Questionnaire**

The questionnaire exhibited excellent reliability, based on Kappa values above .80 for all questions. The time taken for completion ranged from 5 to 10 minutes (mean time, 7.2 minutes; median time, 7.0 minutes). The questionnaire had a Flesch Reading Ease Score of 84.8 and a Flesch-Kincaid Grade Level of 4.2, which correspond to a reading age of approximately 10 years, thus showing satisfactory levels of readability. The Passive Sentences Score formula was 0%.

## **Responses to the Questionnaire**

Questionnaire rank scores at T0 and at T1 are illustrated graphically in Figures 2 and 3, respectively, for both groups.

With respect to Questions #1 and #2, a positive value for the differences between ratings at T0 and at T1 was considered to be indicative of a decrease in patients' opinions regarding the appearance of their smiles or facial profiles. A value of 0 indicated no change, and a negative value suggested an improved opinion. Because only a few improved opinions were found at T1, unchanged and improved opinions were grouped together (Figure 4).

With respect to Questions #3 and #4, a value of 0 for the difference between ratings measured at T0 and at T1 indicated no change in subjects' opinions. A positive difference revealed that subjects were willing to undergo a more comprehensive procedure to change their appearance, and a negative score suggested that they were unwilling. Because only a few subjects were unwilling to undergo a more comprehensive procedure at T1, 0 values and negative scores were grouped together (Figure 4).

# Kruskal-Wallis Equality-of-Populations Rank Test

The *P* values for rank tests are reported in Table 3. For all examined variables, the differences between the two groups were not statistically significant. It was, therefore, possible to assume that subjects in the Study and Control Groups were from the same population.

# **Comparison Between Groups**

No differences were noted between groups with respect to participants' opinions regarding the appearance of their smiles ( $\chi^2$ , 0.34; P > .05) between T0 and

T1. However, a significant difference was found between groups with respect to subjects' opinions regarding the appearance of their facial profiles ( $\chi^2$ , 86.30; *P* < .001) and the types of treatment they would seek to change their smiles ( $\chi^2$ , 15.89; *P* < .001) or facial profiles ( $\chi^2$ , 66.88; *P* < .001) between T0 and T1.

In the Study Group, 11% of subjects were willing to undergo more comprehensive therapies to change their smiles after exposure to photographs of themselves between T0 and T1. Moreover, 50% of patients in the Study Group had a more negative opinion regarding the esthetics of their facial profiles. Accordingly, 45% of subjects were willing to undergo more comprehensive therapies to change their profile appearance. No statistically significant difference was detected in the Control Group between questionnaire ratings at T0 and at T1.

A logistic regression analysis demonstrated that an overjet of >7.1 mm (odds ratio [OR], 0.26; 95% confidence interval [CI], 0.63 to 0.95) and a higher educational level in terms of degrees attained (OR, 0.34; 95% CI, 0.13 to 0.86) were negatively associated, in a statistically significant manner, with a decrease in subjects' opinions between T0 and T1 regarding the appearance of the facial profile. Associations among the remaining variables did not prove to be statistically significant.

# DISCUSSION

Few previous questionnaire-based studies have investigated the self-perception of the attractiveness of the face, especially in the profile view, of individuals seeking orthodontic evaluation.<sup>1</sup> The ability of patients to recognize their profiles from among various silhouettes and photographs<sup>1,8,11,14,18–23</sup> or to reproduce their own profiles had already been assessed.<sup>18,24</sup> Only one recent study investigated whether subjects requiring orthognathic surgery had seen their own facial profiles, and investigators assessed, using a questionnaire, whether subjects were happy with the appearance of their profiles.<sup>25</sup>

In the present study, a large sample of patients who were seeking or were referred for orthodontic evaluation was recruited. Exposure to the view of their pretreatment profiles and smile photographs, as well as discussions with relatives and friends, represented the "treatment variable" in the Study vs Control Group. Care was taken to ensure that the Control Group was matched for age and sex distribution—factors already found to influence subjective esthetic judgment and perception of orthodontic treatment need.<sup>26</sup> Because many adolescents are not fully aware of external motivating factors in the decision to undergo orthodontic treatment, only those individuals aged 18 years or older were included.<sup>1,26,27</sup>



Figure 2. Histograms comparing the questionnaire rank scores of the Study and Control Groups at T0.



Figure 3. Histograms comparing the questionnaire rank scores of the Study and Control Groups at T1.



Figure 4. Histograms comparing changes in subjects' opinions according to questionnaire rank scores (T1-T0; percentages) of the Study vs Control Group.

No differences were found in questionnaire ratings between T0 and T1 for controls, thus implying that subjects not exposed to photographs of themselves did not change their self-perception of dentofacial attractiveness or their willingness to undergo treatment within the assessed period of 30 days. Similarly, no statistically significant difference was noted for either group between T0 and T1 with respect to scores for self-rated happiness with the appearance of their smiles. A statistically significant difference was noted between groups in terms of happiness with the appearance of their facial profiles and willingness to undergo treatment between T0 and T1. The main finding was that subjects exposed to photographs of themselves were significantly less happy with their facial profile appearance than were controls, with 50% of subjects in the Study Group negatively changing their opinions regarding the appearance of their own facial profiles at T1 (Figure 4). This observation is in accordance with previous findings that subjects who

**Table 3.** Kruskal-Wallis Equality-of-Populations Rank Test byGroup (Study vs Control Group)

Variables	P Value
Level of income	.42
Marital status	.62
Level of education	.61
Overjet	.57
Angle class	.71

reported having seen their own faces in profile were less likely to be happy with their profiles.<sup>1,25</sup>

In addition, previous studies have already concluded that patients are somewhat inaccurate in evaluating themselves, particularly regarding their own profiles.<sup>11,18,28</sup> Therefore, it is reasonable to assume that laypeople do not usually see themselves from a lateral view<sup>29</sup> and are thus unfamiliar with their own profiles unless exposed to photographs. Because orthodontic patients may become more conscious of their facial profiles owing to consultations that they receive at the beginning of treatment,<sup>1,11</sup> in the present study care was taken to avoid any esthetic judgments made by the clinician, and, for the same reason, no explanations of the goals of the study were given.

At T1 in the Study Group, 11% of subjects were willing to undergo more comprehensive procedures to change the appearance of their smiles, and 45% were willing to undergo more comprehensive procedures to change the appearance of their facial profiles (Figure 4).

These results confirm that frontal perspectives of the face and the smile are more "familiar" to patients than the lateral view. Because orthodontists place major treatment planning emphasis on the anteroposterior dimension, a difference does exist between professionals' and patients' evaluations. At this point, a question arises as to whether self-perception of the esthetics of the face should be influenced by the orthodontist. It is the authors' opinion that such perception should be considerably guided by orthodontists, who are aware of the extent to which the therapeutic choice, combined with the effects of growth and aging (the so-called fourth dimension), would influence the overall dentofacial esthetic outcome. While discussing orthodontic diagnosis and treatment planning, patients' exposure to photographs proved to be a valuable tool in making the perceptions of patients match more closely those of orthodontists. This alignment of perceptions would result in more realistic motivations for and expectations of treatment on the part of patients.

#### CONCLUSION

- Laypeople are not generally aware of their facial profiles unless exposed to photographs.
- Exposure to pretreatment facial photographs would reduce the discrepancy between patients' actual and perceived levels of facial attractiveness, thus making orthodontists' and patients' visual emphasis on dentofacial esthetics more similar to one another.

#### REFERENCES

- 1. Tufekci E, Jahangiri A, Lindauer SJ. Perception of profile among laypeople, dental students and orthodontic patients. *Angle Orthod.* 2008;78:983–987.
- 2. Cunningham SJ, Hunt NP, Feinmann C. Psychological aspects of orthognathic surgery: a review of the literature. *Int J Adult Orthodon Orthognath Surg.* 1995;10:159–172.
- 3. Soh J, Chew MT, Chan YH. Perceptions of dental esthetics of Asian orthodontists and laypersons. *Am J Orthod Dentofacial Orthop.* 2006;130:170–176.
- Schlosser JB, Preston CB, Lampasso J. The effects of computer-aided anteroposterior maxillary incisor movement on ratings of facial attractiveness. *Am J Orthod Dentofacial Orthop.* 2005;127:17–24.
- 5. Shafiee R, Korn EL, Pearson H, Boyd RL, Baumrind S. Evaluation of facial attractiveness from end-of-treatment facial photographs. *Am J Orthod Dentofacial Orthop.* 2008; 133:500–508.
- Soh J, Chew MT, Wong HB. A comparative assessment of the perception of Chinese facial profile esthetics. *Am J Orthod Dentofacial Orthop.* 2005;127:692–699.
- Siqueira DF, Sousa MV, Carvalho PE, Valle-Corotti KM. The importance of the facial profile in orthodontic diagnosis and treatment planning: a patient report. *World J Orthod.* 2009; 10:361–370.
- 8. Foster EJ. Profile preferences among diversified groups. *Angle Orthod*. 1973;43:34–40.
- Maple JR, Vig KW, Beck FM, Larsen PE, Shanker S. A comparison of providers' and consumers' perceptions of facialprofile attractiveness. *Am J Orthod Dentofacial Orthop.* 2005;128:690–696.
- Isıksal E, Hazar S, Akyalçın S. Smile esthetics: perception and comparison of treated and untreated smiles. *Am J Orthod Dentofacial Orthop.* 2006;129:8–16.
- 11. McKoy-White J, Evans CA, Viana G, Anderson NK, Giddon DB. Facial profile preferences of black women before and

after orthodontic treatment. Am J Orthod Dentofacial Orthop. 2006;129:17–23.

- Gracco A, Cozzani M, D'Elia L, Manfrini M, Peverada C, Siciliani G. The smile buccal corridors: aesthetic value for dentists and laypersons. *Prog Orthod*. 2006;7:56–65.
- 13. Park YS, Evans CA, Viana G, Anderson NK, Giddon DB. Profile preferences of Korean American orthodontic patients and orthodontists. *World J Orthod.* 2006;7:286–292.
- 14. Coleman GG, Lindauer SJ, Tüfekçi E, Shroff B, Best AM. Influence of chin prominence on esthetic lip profile preferences. *Am J Orthod Dentofacial Orthop.* 2007;132:36–42.
- Miner RM, Anderson NK, Evans CA, Giddon DB. The perception of children's computer-imaged facial profiles by patients, mothers and clinicians. *Angle Orthod.* 2007;77: 1034–1039.
- Schabel BJ, McNamara JA Jr, Franchi L, Baccetti T. Q-sort assessment vs visual analog scale in the evaluation of smile esthetics. *Am J Orthod Dentofacial Orthop.* 2009;135: S61–S71.
- Schabel BJ, Franchi L, Baccetti T, McNamara JA Jr. Subjective vs objective evaluations of smile esthetics. *Am J Orthod Dentofacial Orthop.* 2009;135:S72–79.
- Hershon LE, Giddon DB. Determinants of facial profile selfperception. Am J Orthod Dentofacial Orthop. 1980;78:279– 295.
- Arpino VJ, Giddon DB, BeGole EA, Evans CA. Presurgical profile preferences of patients and clinicians. *Am J Orthod Dentofacial Orthop.* 1998;114:631–637.
- Hall D, Taylor RW, Jacobson A, Sadowsky PL, Bartolucci A. The perception of optimal profile in African Americans versus white Americans as assessed by orthodontists and the lay public. *Am J Orthod Dentofacial Orthop.* 2000;118: 514–525.
- 21. Orsini MG, Huang GJ, Kiyak HA, et al. Methods to evaluate profile preferences for the anteroposterior position of the mandible. *Am J Orthod Dentofacial Orthop.* 2006;130: 283–291.
- Xu T, Korn EH, Liu Y, et al. Facial attractiveness: ranking of end-of-treatment facial photographs by pairs of Chinese and US orthodontists. *Am J Orthod Dentofacial Orthop.* 2008; 134:74–84.
- Liu Y, Korn EL, Oh HS, Pearson H, Xu T, Baumrind S. Comparison of Chinese and US orthodontists' averaged evaluations of "facial attractiveness" from end-of-treatment facial photographs. *Am J Orthod Dentofacial Orthop.* 2009; 135:621–634.
- Giddon DB, Hershon LE, Lennartsson B. Discrepancy between objective and subjective profile measures. *Scand J Dent Res.* 1974;82:527–535.
- Johnston C, Hunt O, Burden D, Stevenson M, Hepper P. Self-perception of dentofacial attractiveness among patients requiring orthognathic surgery. *Angle Orthod.* 2010;80: 361–366.
- Wedrychowska-Szulc B, Syryńska M. Patient and parent motivation for orthodontic treatment—a questionnaire study. *Eur J Orthod.* 2010;32:447–452.
- Trulsson U, Strandmark M, Mohlin B, Berggren U. A qualitative study of teenagers' decision to undergo orthodontic treatment with fixed appliance. *J Orthod.* 2002;29: 197–204.
- Pitt EJ, Korabik K. The relationship between self-concept and profile self-perception. *Am J Orthod Dentofacial Orthop*. 1977;72:459–460.
- 29. Soh J, Chew MT, Wong HB. An Asian community's perspective on facial profile attractiveness. *Community Dent Oral Epidemiol.* 2007;35:18–24.