A comparison of the effects of Forsus appliances with and without temporary anchorage devices for skeletal Class II malocclusion: *A systematic review and meta-analysis*

Lu Liu^a; Qi Zhan^b; Jing Zhou^a; Qianyun Kuang^a; Xinyu Yan^a; Xiaoqi Zhang^a; Yue Shan^a; Wenli Lai^c; Hu Long^d

ABSTRACT

Objectives: To compare the effects of Forsus appliances with and without temporary anchorage devices (TADs) for patients with skeletal Class II malocclusion.

Materials and Methods: Through a predefined search strategy, electronic searching was conducted in PubMed, Embase, Web of Science, CENTRAL, ProQuest Dissertations & Theses, and SIGLE with no language restrictions. Eligible study selection, data extraction, and evaluation of risk of bias (Cochrane Collaboration tool) were conducted by two authors independently and in duplicate. Any disagreement was solved by discussion or judged by a third reviewer. Statistical pooling, sensitivity analysis, subgroup analysis, and assessment of small-study effects were conducted by using Comprehensive Meta-Analysis and Stata 12.0. Heterogeneity was analyzed for different types of study designs, TADs, and radiographic examinations.

Results: Electronic search yielded a total of 256 studies after removing duplicates. Among them, six studies were finally included. All articles were of high quality. The pooled mean differences were -0.27 (95% confidence interval [CI]: -0.59, 0.05) for SNA, 0.58 (95% CI: -0.07, 1.23) for SNB, -0.86 (95% CI: -1.74, -0.03) for ANB, 1.63 (95% CI: 0.46, 2.80) for Co-Po, 0.75 (95% CI: 0.28, 1.23) for SN-MP, -7.56 (95% CI: -11.37, -3.76) for L1-MP, 0.47 (95% CI: -0.98, 1.91) for overjet, 0.39 (95% CI: -0.57, 1.35) for overbite, -1.84 (95% CI: -5.15, 1.47) for SN-OP, and 4.97 (95% CI: -1.22, 11.17) for nasolabial angle.

Conclusions: TADs (especially miniplates) were able to eliminate dental adverse effects of Forsus appliances for correction of skeletal Class II malocclusion. (*Angle Orthod.* 2021;91:255–266.)

KEY WORDS: Class II malocclusion; Forsus; Temporary anchorage devices; Systematic review; Meta-analysis

INTRODUCTION

Class II malocclusion is a constellation of several types of malocclusion. Patients with Class II malocclusion can exhibit maxillary protrusion, mandibular retrusion, or their combinations, together with abnormal dental relationship problems and facial esthetic disorders.¹ In particular, mandibular retrusion, rather than maxillary protrusion, is the main etiologic factor for

^b Undergraduate student, West China School of Stomatology, Sichuan University, Sichuan, China.

(e-mail: hlong@scu.edu.cn)

Accepted: October 2020. Submitted: May 2020.

^a Graduate student, State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases & Department of Orthodontics, West China Hospital of Stomatology, Sichuan University, Sichuan, China.

[°] Professor, State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases & Department of Orthodontics, West China Hospital of Stomatology, Sichuan University, Sichuan, China.

^d Associate Professor, State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases & Department of Orthodontics, West China Hospital of Stomatology, Sichuan University, Sichuan, China.

Corresponding author: Dr Hu Long, State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases & Department of Orthodontics, West China Hospital of Stomatology, Sichuan University; No.14, Section 3, Ren Min South Road, Chengdu, Sichuan 610041, China

Published Online: December 30, 2020

^{© 2021} by The EH Angle Education and Research Foundation, Inc.

Step	PubMed Search Strategy
1	"Malocclusion, Angle Class II" [mesh] OR Angle Class II OR Class II, Angle OR Class II OR Distocclusion* OR Distal occlusion*
2	Forsus OR functional appliance OR orthopedic OR mandibular advancement
3	Dental implant OR Micro-implant OR Microimplant OR Micro implant OR Mini-implant OR Mini implant OR Mini-screw OR Miniscrew OR Mini screw OR Mini-plate OR Mini plate OR Mini plate OR Skeletal anchorage OR Bone anchorage OR Bone screw
4	1 AND 2 AND 3

Table 1. Search Strategy for PubMed

Class II malocclusion² and the most common characteristic of Class II malocclusion.³ Therefore, orthodontic treatments promoting mandibular growth or advancement are indicated for growing patients with mandibular retrusion.

A variety of orthodontic appliances have been designed for mandibular retrusion, including fixed functional and removable functional appliances. They achieve Class II correction by inducing mandibular lengthening sagittally and vertically by stimulating condylar growth.4 While the treatment effects of removable functional appliances are limited by poor patient cooperation, fixed functional appliances (FFAs) are more advantageous and are able to achieve clinically significant mandibular elongation.⁵ Among various FFAs, Forsus Fatigue Resistant Device (FFRD; 3M Unitek, Monrovia, Calif)⁶ is an example of a hybrid FFA, which is used for treatment of mandibular retrusion in growing patients. However, it has been suggested that Forsus could induce mandibular lengthening at the expense of dentoalveolar compensations. Forsus exerts anteriorly directed force on the mandibular dentition, and this may result in proclination of the mandibular incisors, jeopardizing the long-term stability of treatment outcomes.7

To minimize the adverse effects of Forsus, a combination of Forsus and temporary anchorage devices (TADs) has been claimed to preserve dental anchorage and avoid adverse dental effects.^{8–12} However, controversy arose regarding whether Forsus with and without skeletal anchorage produced similar treatment results.^{9,13} These conflicting results are likely attributed to different types of skeletal anchorage and different study designs. To date, no conclusion has been made whether Forsus with TADs are superior to Forsus only. The aims of this study were to compare the effects of Forsus with and without TADs for patients with Class II mandibular retrusion.

MATERIALS AND METHODS

Protocol and Registration

The protocol for this systematic review was registered in the International Prospective Register of Systematic Reviews (PROSPERO; http://www.crd.york.ac.uk/ prospero; registration number CRD42020140721).

Inclusion Criteria

Growing patients diagnosed with skeletal Class II base with mandibular retrusion were included. Interventions were Forsus and fixed appliances with or without TADs (eg, miniscrews and miniplates) for mandibular advancement. Both randomized controlled trials (RCTs) and prospective controlled clinical trials (CCTs) were eligible.

Exclusion Criteria

Patients with severe orofacial anomalies (eg, cleft lip), dental pathology (eg, cyst), and medical conditions (eg, osteoporosis) were excluded.

Information Sources, Search Strategy, and Study Selection

The databases searched included those of PubMed, Embase, Web of Science, Cochrane Central Register of Controlled Trials (CENTRAL), ProQuest Dissertations & Theses, and the gray literature database of SIGLE. Specifically, the electronic search parameters were articles from January 1980 to April 2020 with no language restrictions. The specific search strategy of PubMed is presented in Table 1. Titles and abstracts were assessed according to the eligibility criteria. The selected articles were evaluated after reading full texts, and a final selection was determined. Searching and assessment of studies was performed independently and in duplicate by two reviewer authors (Miss Liu and Miss Zhan), and disagreements were judged by a third reviewer (Dr Long).

Data Extraction

Results regarding study design, demographic data (age, sample size, grouping, and skeletal age), detailed description (interventions, treatment duration, and comparator), and outcomes (skeletal, dentoalveolar, and soft-tissue measurements from anteroposterior, and vertical direction) were all extracted.

Study Outcomes

Study outcomes were skeletal indices (SNA, SNB, ANB, Co-Po, SN-MP), dental indices (L1-MP, overjet, overbite, SN-OP), and soft-tissue indices (nasolabial

angle). Primary outcomes were ANB, SN-MP, and L1-MP, and secondary outcomes were SNA, SNB, Co-Po, overjet, overbite, SN-OP, and nasolabial angle.

Risk of Bias and Quality Assessment

The evaluation of risk of bias for the included studies was performed independently and in duplicate by two review authors (Miss Liu and Miss Zhan) according to the Cochrane tool for assessing risk of bias in randomized trials (RoB 2.0 tool)¹⁴ and in nonrandomized trials (ROBINS-I tool).¹⁵ For each bias domain, a judgment score was given following the recommendations of the Cochrane Handbook for Systematic Reviews of Interventions 5.1.0 Any disagreement was solved by discussion or judged by a third reviewer (Dr Long). Additionally, the Grading of Recommendation Assessment, Development and Evaluation (GRADE) system of rating quality of evidence was used to appraise the quality of each outcome in this review.¹⁶⁻²¹

Meta-analysis

Original outcome data were subject to statistical pooling through random effect models. The criteria of data pooling were determined a priori on the basis of the comparability of study design, patient type, treatments, outcomes, and risk of bias. Mean difference (MD) was used for statistical pooling for continuous data. Heterogeneity across studies was assessed through the I² statistic, and an I² statistic greater than 50% was considered as substantial heterogeneity.

If substantial heterogeneity existed, a subgroup analysis on different study designs, types of TADs, and radiographic modalities was executed to explore the potential heterogeneity. In addition, sensitivity analysis for studies with small sample sizes was conducted to evaluate the robustness of the pooled results in the meta-analysis. Cumulative meta-analysis was performed to determine the chronological changes of the pooled results. The Egger test and Begg test were used to assess publication bias or small study effect.

All the meta-analyses were performed in Comprehensive Meta-Analysis (version 2.2.064, Biostat, Englewood, NJ) and Stata 12.0 (Stata Corp, College Station, Tex).

RESULTS

Description of Studies

Initially, 256 articles from the database were identified and 248 were excluded as irrelevant. The remaining eight articles were further assessed for



Figure 1. PRISMA flow diagram for studies retrieved through the search and selection processes.

eligibility, and six studies (three RCTs and three prospective CCTs) were finally included in this systematic review.⁸⁻¹³ The procedures of electronic searching and selection are shown in Figure 1. Sample sizes ranged from 8 to 16 in different groups, with ages between 12 and 14 years old. The active Forsus treatment durations ranged from 4.86 months to 10.45 months. The details of each study are presented in Table 2. All six articles⁸⁻¹³ were high quality. Risk of bias is presented in Table 3.

Description of Interventions

Forsus appliances and fixed appliances were used in all the included studies. Two types of TADs were used among the included studies: miniscrew and miniplate. Among the included studies, three studies^{8,9,13} used miniscrews and the other three studies^{10–12} used miniplates. A total of 169 participants were included in this systematic review: 84 received Forsus+TADs and 85 received Forsus only. Among the 84 participants receiving Forsus+TADs, miniscrews were used in 46 participants, and the remaining 38 received miniplates.

Description of Outcomes

Among the proposed study outcomes, 10 outcomes were investigated in the included studies: skeletal outcomes (SNA, SNB, ANB, Co-Po, and SN-MP), dental outcomes (L1-MP, overjet, overbite, and SN-OP), and soft-tissue outcome (nasolabial angle). Outcomes at the active treatment effect period (changes from before Forsus treatment to after Forsus treatment) were studied in this meta-analysis. The GRADE assessment for quality of evidence for each outcome is shown in Table 4. Unfortunately, the quality of evidence of the outcomes in this meta-analysis was determined to be very low.

Study	Design	Participants	Skeletal Class II	Skeletal Maturation	Treatment Duration (mo)	Fixed Appliance	TAD	Radiograph Analysis
Aslan et al. 2014 [°]	Prospective CCT	TAD: n = 16, age = 13.68 ± 1.09 No TAD: n = 17, age = 14.64 ± 1.56	Class II molar relationship with mandibular retrognathia	N/A	TAD: 6.5 ≟ 1.97 No TAD: 5.5 ≟ 1.80	0.018-inch slot Roth brackets	Miniscrew	Lateral cephalometric
Eissa et al. 2017'³	RCT	TAD: n = 15, age = 12.52 ± 1.12 No TAD: n = 14, age = 12.76 ± 1.0	$\begin{array}{l} ANB > 4.5^{\circ} \\ SNB < 76^{\circ} \end{array}$	Stage 2-4 (CVM)	TAD: 6.42 ± 1.04 No TAD: 6.06 ± 0.76	0.022-inch slot MBT brackets	Miniscrew	Lateral cephalometric
Elkordy et al. 2019 ¹⁰	RCT	TAD: n = 15, age = 12.06 ± 0.79 No TAD: n = 15, age = 12.54 ± 0.90	SNB $< 76^{\circ}$	Stage 3-4 (CVM)	TAD: 9.42 ± 0.98 No TAD: 6.23 ± 1.61	0.022-inch slot MBT brackets	Miniplate	CBCT
Elkordy et al. 2016°	RCT	TAD: n = 15, age = 13.07 ± 1.41 No TAD: n = 16, age = 13.25 ± 1.12	SNB $< 76^{\circ}$	Stage G–H (MP3)	TAD: 5.34 ± 1.29 No TAD: 4.86 ± 1.32	0.022-inch slot MBT brackets	Miniscrew	CBCT
Gandedkar et al. 2019 ¹¹	Prospective CCT	TAD: n = 8, age = 12.96 ± 0.38 No TAD: n = 8, age = 13.11 ± 0.38	SNA 84 \pm 2° SNB 76 \pm 2° ANB > 4 \pm 1°	15% prepubertal 70% pubertal 15% postpubertal	TAD: 10.45 ± 0.16 No TAD: 7.59 ± 0.32	0.022-inch slot brackets	Miniplate	CBCT
Turkkahraman et al. 2016¹²	Prospective CCT	TAD: n = 15, age = 12.77 ± 1.24 No TAD: n = 15, age = 13.26 ± 0.82	Angle Class II molar relationship; convex profile with mandibular deficiency	N/A	TAD: 9.40 ± 2.25 No TAD: 9.46 ± 0.81	0.018-inch slot Roth brackets	Miniplate	Lateral cephalom-etric
^a CBCT indica vertebrae matura	tes cone-beam (tion; MP3, midd	computed tomography; Co	CT, controlled clinical trial; N/A, inger; MBT, McLaughlin, Benn	, not applicable; RCT, ett and Trerisi.	randomized contro	olled trial; TAD, tempo	orary anchoraç	e device; CVM, cervical

Table 2. General Information Provided in the Included Studies $^{\rm a}.$

Table 3. Risk of Bias of the Included Studies

Study	Item 1	Item 2	Item 3	Item 4	Item 5	Item 6	Item 7	Quality
ROB 2.0ª								
Elkordy et al. 2016°	Low	Low	Low	Low	Low	Low		High
Eissa et al. 201713	Low	Low	Unclear	Low	Low	Unclear		High
Elkordy et al. 201910	Low	Low	Unclear	Low	Low	Unclear		High
ROBINS-I ^b								•
Aslan et al. 2014 ⁸	Unclear	Low	Low	Low	Low	Unclear	Low	High
Turkkahraman et al. 201612	Unclear	Low	Low	Low	Low	Unclear	Low	High
Gandedkar et al. 201911	Unclear	Low	Low	Low	Low	Unclear	Low	High

^a ROB 2.0: item 1, bias arising from the randomization process; item 2, bias due to deviations from intended interventions; item 3, bias due to missing outcome data; item 4, bias in measurement of the outcome; item 5, bias in selection of the reported result; and item 6, overall bias. Scoring rules: low indicates a score of 2; unclear, a score of 1; and high, a score of 0. Quality was categorized as low (score 1–4), medium (score 5–8), or high (score 9–12).

^b ROBINS-I: item 1, bias due to confounding; item 2, bias in selection of participants into the study; item 3, bias in classification of interventions; item 4, bias due to deviations from intended interventions; item 5, bias due to missing data; item 6, bias in measurement of outcomes; and item 7, bias in selection of the reported result. Scoring rules: low indicates a score of 2; unclear, a score of 1; and high, a score of 0. Quality was categorized as low (score 1–5), medium (score 6–10), or high (score 11–14).

Meta-analysis

For skeletal outcomes (Figure 2), six studies^{8–13} investigated SNA, SNB, and ANB angle, which had pooled MDs of -0.27° (95% confidence interval [CI]: -0.59, 0.05), 0.58° (95% CI: -0.07, 1.23), and -0.86° (95% CI: -1.74, -0.03), respectively. Five studies^{8,10–13} investigated Co-Po, which had a pooled MD of 1.63 mm (95% CI: 0.46, 2.80). Six studies^{8–13} investigated SN-MP, which had a pooled MD of 0.75° (95% CI: 0.28, 1.23).

For dental outcomes (Figure 3), four studies^{8,10-12} investigated L1-MP, which had a pooled MD of -7.56° (95% CI: -11.37, -3.76). Four studies^{8,11-13} investigated overjet and overbite, which had pooled MDs of 0.47 mm (95% CI: -0.98, 1.91) and 0.39 mm (95% CI: -0.57, 1.35), respectively. Two studies^{8,12} investigated

Table 4. GRADE Assessment for Quality of Evidence

SN-OP, which had a pooled MD of -1.84° (95% CI: -5.15, 1.47).

For soft-tissue outcome (Figure 4), one study¹³ investigated nasolabial angle, which was 4.97° (95% CI: -1.22, 11.17).

Sensitivity Analysis and Subgroup Analysis

Due to the small sample size in Gandedkar et al.,¹¹ sensitivity analysis was performed by excluding this study from the meta-analysis. Stable results were found except for Co-Po (MD = 1.34; 95% CI: $-0.36 \sim 3.04$).

Subgroup analysis was performed since different study designs, TADs, and radiographic modalities could influence the results of the meta-analysis (Table 5).

				Downgrade	3			Overall
Index	Study Design	Limitations	Inconsistency	Indirectness	Imprecision	Publication Bias	Upgrade	Quality
SNA	3 RCTs	Serious	Serious	None	Serious	None	None	Very low
	3 prospective CCTs							
SNB	3 RCTs	Serious	Very serious	None	Very serious	None	None	Very low
	3 prospective CCTs							
ANB	3 RCTs	Serious	Very serious	None	Very serious	Likely	None	Very low
	3 prospective CCTs							
Co-Po	2 RCTs	Serious	Very serious	None	Very serious	Likely	None	Very low
	3 prospective CCTs							
SN-MP	3 RCTs	Serious	Serious	None	Serious	None	None	Very low
	3 prospective CCTs							
L1-MP	1 RCT	Serious	Serious	None	Very serious	None	None	Very low
	3 prospective CCTs							
Overjet	1RCT	Serious	Very serious	None	Very serious	None	None	Very low
	3 prospective CCTs							
Overbite	1 RCT	Serious	Very serious	None	Serious	None	None	Very low
	3 prospective CCTs							
SN-OP	2 prospective CCTs	Serious	Serious	None	Very serious	None	None	Very low
Nasolabial angle	1 RCT	None	None	None	Very serious	None	None	Very low

^a CCT controlled clinical trial; RCT, randomized controlled trial.



Figure 2. Forest plots of pooled MD of skeletal outcomes (SNA, SNB, ANB, Co-Po, and SN-MP) for the TAD-FFAs group versus control group.



No TADs TADs

Figure 3. Forest plots of pooled MD of dental outcomes (L1-MP, overjet, overbite, and SN-OP) for the TAD-FFAs group versus control group.



Figure 4. Forest plots of pooled MD of soft-tissue outcome (nasolabial angle) for the TAD-FFAs group versus control group.

In terms of study designs, included studies were divided into two subgroups: CCTs^{8,11,12} and RCTs.^{9,10,13} Significant changes from original estimates were detected for ANB, Co-Po, and SN-MP.

For different types of TADs, two subgroups were analyzed: miniscrew^{8,9,13} and miniplate.^{10–12} The subgroup analysis revealed that ANB and Co-Po differed from original results for the subgroup of miniscrew, and ANB and SN-OP were different from original results for the miniplate subgroup.

For different modalities of radiography, the subgroup of lateral cephalometric^{8,12,13} and the subgroup of conebeam computed tomography (CBCT)^{9–11} were available. ANB, Co-Po in the subgroup of lateral cephalometry, and SNA, ANB in the subgroup of CBCT were different from original results.

Cumulative Meta-analysis

Some of the important results of cumulative analysis are displayed in Figure 5. The results of SNA, SNB, L1-MP, overjet, SN-OP, and nasolabial angle were stable, while unstable results were found for ANB, Co-Po, SN-MP, and overbite.

Assessment of Publication Bias

The assessment of publication bias is shown in Table 6. Because of the limited number of studies that analyzed SN-OP and nasolabial angle, assessment of publication bias was not performed for these indices. For other indices, the Egger test and Begg test revealed no evidence of publication bias except for Co-Po.

DISCUSSION

In this systematic review, skeletal, dental, and softtissue outcomes were examined to compare the effectiveness of mandibular advancement with and without TADs. Despite heterogeneity among the studies included, subgroup analysis was performed to explore heterogeneity. Neither the Begg test nor the Egger test detected any evidence of publication bias except for Co-Po. In addition, although sensitivity and subgroup analyses revealed significant changes from original estimates, these significant changes could be explained. Thus, the results in this meta-analysis were robust.

As shown in Figure 2, the skeletal outcomes included SNA, SNB, ANB, Co-Po, and SN-MP. These skeletal outcomes could be subdivided into sagittal outcomes (SNA, SNB, ANB, and Co-Po) and vertical outcomes (SN-MP). The pooled SNA was -0.27° (95% CI: -0.59, 0.05), and sensitivity analysis revealed robust pooled results regarding SNA, indicating that the use of TADs failed to achieve additional maxillary restriction. However, the subgroup analysis on different radiographic modalities revealed that TADs could reduce SNA for the CBCT subgroup but not for the lateral cephalometric radiography subgroup (Table 5). This could be explained by the improved visualization of CBCT over lateral cephalometrics.²² Despite this, the reduction of SNA with the use of TADs for the CBCT subgroup (MD: –0.49; 95% CI: –0.90 \sim –0.07) was of no clinical significance. Therefore, it is suggested that the additional use of TADs had no benefits in maxillary restriction compared with Forsus alone.

With regard to ANB, the meta-analysis revealed a significant difference between TADs and no TADs. Cumulative meta-analysis revealed no significant difference between TADs and no TADs before 2019, but the pooled ANB became significantly different afterward. This might be attributed to the use of miniplates. However, subgroup analysis showed no significantly different results in different types of study designs, TADs, or radiography, which indicated that the pooled result was unstable. This was in agreement with previous studies.^{8,9,12,13} Therefore, due to the unstable results for ANB, it cannot be determined whether Forsus + TADs could bring about more skeletal correction over Forsus only.



Figure 5. Cumulative meta-analysis of ANB, Co-Po, SN-MP, and L1-MP for the TAD-FFAs group versus control group.

		Study I	Design	17	ADs	Radioc	jraphy
Index	Original result	Prospective CCT	RCT	Miniscrew	Miniplate	Lateral Cephalometric	CBCT
SNA	-0.27 (-0.59, 0.05)	-0.21 (-0.94, 0.51)	-0.28 (-0.56, 0.00)	-0.23 (-0.47, 0.01)	-0.32 (-1.16, 0.53)	-0.05 (-0.54, 0.44)	-0.49 (-0.90, - 0.07)
SNB	0.58 (-0.07, 1.23)	0.83 (-0.84, 2.50)	0.31 (-0.03, 0.65)	0.15 (-0.13, 0.42)	1.13 (-0.30, 2.56)	0.00 (-0.30, 0.29)	1.19 (-0.08, 2.47)
ANB	-0.86 (-1.74, -0.03)	-1.20 (-3.14, 0.74)	-0.51 (-1.07, 0.04)	-0.22(-0.48, 0.03)	-1.58 (-3.36, 0.21)	-0.24(-0.56, 0.09)	-1.56 (-3.42, 0.29)
Co-Po	1.63 (0.46, 2.80)	1.36 (0.06, 2.65)	1.94 (-0.77, 4.65)	0.25 (-0.92, 1.43)	2.38 (1.32, 3.43)	0.69 (-0.17, 1.55)	2.85 (2.07, 3.64)
SN-MP	0.75 (0.28, 1.23)	0.59 (0.08, 1.09)	0.92 (-0.01, 1.84)	0.52 (0.06, 0.99)	1.06 (0.11, 2.02)	0.62 (0.08, 1.17)	0.91 (0.04, 1.78)
L1-MP	-7.56 (-11.37, -3.76)	-6.43 (-10.04, -2.81)	-10.67 (-13.35, -8.00)	-5.68 (-8.73, -2.63)	-8.23 (-13.46, -3.00)	-8.01 (-12.74, -3.28)	-7.21 (-13.74, -0.68)
Overjet	0.47 (-0.98, 1.91)	1.03 (-1.04, 3.10)	-1.08 (-2.19, 0.03)	-0.75 (-1.62, 0.13)	1.68 (-1.92, 5.29)	0.74 (-1.99, 3.47)	-0.08 (-0.33, 0.17)
Overbite	0.39 (-0.57, 1.35)	0.81 (-0.43, 2.04)	-0.90 (-2.05, 0.25)	0.15 (-1.86, 2.16)	0.71 (-1.18, 2.60)	0.66 (-0.88, 2.21)	-0.13 (-0.31, 0.05)
SN-OP	1.84 (-5.15, 1.47)	1.84 (-5.15, 1.47)	I	-0.12 (-2.36, 2.12)	-3.50 (-5.58, -1.42)	1.84 (-5.15, 1.47)	
Naslabial angle	4.97 (-1.22, 11.17)	I	4.97 (-1.22, 11.17)	4.97 (-1.22, 11.17)	I	4.97 (-1.22, 11.17)	I
^a CBCT indicat	es cone-beam compute	ed tomography; CCT, con	trolled clinical trial; N/A, r	not applicable; RCT, ra	ndomized controlled trial;	TAD, temporary anch	5

Table 6. Assessment of Publication Bias

Index	Begg Test	Egger Test
SNA	0.25966	0.57216
SNB	0.25966	0.44270
ANB	0.06029	0.28487
Co-Po	0.08641	0.00250
SN-MP	0.70711	0.55672
L1-MP	0.73410	0.13565
Overjet	0.30818	0.65692
Overbite	0.73410	0.43573
SN-OP	_	-
Nasolabial angle	-	_

^a Bold values indicate significant difference from the original estimates (P < 0.05).

The pooled results regarding mandibular position were conflicting: the pooled SNB was indicative of no difference between TADs and no TADs, while the pooled Co-Po revealed a significant difference. For SNB, both sensitivity and subgroup analyses were indicative of the robustness of the pooled results (Table 5). However, sensitivity analysis on Co-Po revealed an unstable pooled result (MD: 1.34; 95% CI: -0.36 \sim 3.04) after excluding Gandedkar et al.¹¹ Then, subgroup analysis revealed that the results in different types of study design, TAD, and radiography were also unstable. This conflicting finding could be attributed to the instability of B point, which is not only a skeletal reference but also a dentoalveolar reference. Specifically, it has been shown that lower incisor proclination could cause an advancement of B point.²³ Additionally, Forsus alone could lead to lower incisor proclination compared with the combined use of Forsus and TADs.^{8,10–12} It is suggested that the superiority of mandibular advancement by the combined use of Forsus and TADs could be offset by more advancement of B point (incisor proclination) by Forsus alone. Thus, due to the unstable pooled result, current evidence is unable to verify the clinical effect of TADs for mandibular advancement.

For vertical changes, a significant difference was found for SN-MP, indicating that Forsus with TADs could increase the mandibular plane angle compared with Forsus without TADs, which was in agreement with previous studies.^{9,10,12} The results of sensitivity analysis and subgroup analysis were mostly stable. From a biomechanical standpoint, Forsus alone caused lower incisor proclination, while Forsus with TADs avoided this side effect. As a result, the lower incisors were more upright in Forsus + TADs than in Forsus alone, which resulted in anterior occlusal interference and subsequent clockwise rotation of the mandibular plane in Forsus + TADs group. Therefore, it can be verified that Forsus with TADs can increase the mandibular angle compared with Forsus alone. The dental outcomes including L1-MP, Overjet, Overbite and SN-OP are presented in Figure 3. There were significant results for L1-MP (MD = -7.56° , 95% CI: $-11.37 \sim -3.76$), while there were no significant results for overjet, overbite and SN-OP.

Conventional Forsus exerts anterior forces on the mandibular dentition and the side effect is embodied in proclination of the lower incisors.7 The meta-analysis of L1-MP showed significant reduction of lower incisor proclination with the combination of TADs, which was consistent with previous studies.8,10-12 Both sensitivity and subgroup analyses were robust regarding the pooled results. In addition, cumulative meta-analysis showed that the pooled result was consistent with the original estimate. In particular, it is noteworthy that more reduction in incisor proclination was achieved by miniplates than miniscrews (Table 5). This could be explained by the direct (miniplate) vs indirect (miniscrew) anchorage preservation between the two modalities. Thus, it is suggested that TADs (both miniplates and miniscrews) with Forsus are able to prevent mandibular incisors from proclining compared with Forsus alone, with miniplates being more effective.

No significant results were detected for overjet and overbite, which can be explained by the requirements of study design: edge-to-edge relationship should be achieved at finish. Sensitivity and subgroup analyses were indicative of the robustness of the pooled results, so it is suggested that the additional use of TADs had no effect on overjet and overbite.

SN-OP was similar between TADs and no TADs $(MD = 1.84; 95\% CI: -5.15 \sim 1.47)$. Interestingly, the subgroup analysis on miniplates detected a significant decrease of occlusal plane angle (MD = -3.50; 95% CI: $-5.58 \sim -1.42$), while that on miniscrews did not (MD = -0.12; 95% CI: $-2.36 \sim 2.12$). It was well documented that Forsus could exert intrusive forces on mandibular incisors and produce clockwise rotations of occlusal planes.²⁴ The adjunctive use of miniplates could negate the intrusive forces of Forsus on the mandibular incisors and prevent the occlusal planes from undergoing clockwise rotation, which could produce an anticlockwise rotation of occlusal planes. This is beneficial for patients with Class II malocclusion (especially for hyperdivergent cases) needing mandibular advancement since occlusal planes are the sliding planes for mandibular advancement, and a flatter occlusal plane offers a more horizontal sliding plane for mandibular advancement. In contrast, the adjunctive use of miniscrews did not mitigate the intrusive forces on the lower incisors by Forsus and, thus, had no effect on the occlusal plane.

For soft-tissue outcomes (Figure 4), only nasolabial angle was examined among the studies included. The

results revealed that the nasolabial angle was not significantly different between Forsus with and without TAD treatment. However, due to insufficient evidence, this finding should be interpreted with caution.

Limitations

In this meta-analysis, most of the studies included had a moderate sample size, but Gandedkar et al.¹¹ had a small sample size (eight participants per group). The small sample size could cause strongly positive results and affect the pooled results. In addition, the sample sizes for quality assessment of the outcomes were small for most of the indices, and imprecision of the outcomes could be serious.

Different study designs might have had a confounding influence on the results of the outcomes. Thus, future RCTs with uniform study designs are warranted.

The included studies only examined short-term effects on mandibular advancement and dental side effects. Therefore, results of this meta-analysis cannot support conclusions on the long-term effects of Forsus with or without TADs.

CONCLUSIONS

On the basis of this review, we drew the following conclusions. However, due to the low quality of study outcomes, conclusions should be interpreted with caution:

- Current evidence did not verify the additional sagittal skeletal effects of TADs in conjunction with Forsus appliances.
- Forsus with TADs resulted in mandibular clockwise (opening) rotation compared with those without TADs.
- Forsus with TADs could avoid lower incisor proclination; miniplates were more effective than miniscrews.
- Miniplates could, while miniscrews could not, prevent the occlusal plane from being steepened.
- The effects of TADs in conjunction with Forsus could not be determined for soft tissues.

ACKNOWLEDGMENTS

This study was supported by National Natural Science Foundation of China (No. 82071147, 81571004, and 81500884).

REFERENCES

- Arici S, Akan H, Yakubov K, Arici N. Effects of fixed functional appliance treatment on the temporomandibular joint. *Am J Orthod Dentofacial Orthop.* 2008;133:809–814.
- McNamara JA Jr. Components of class II malocclusion in children 8-10 years of age. *Angle Orthod*. 1981;51:177–202.

- 3. Perillo L, Padricelli G, Isola G, et al. Class II malocclusion division 1: a new classification method by cephalometric analysis. *Eur J Paediatr Dent*. 2012;13:192–196.
- Vaid NR, Doshi VM, Vandekar M. Class II treatment with functional appliances: a meta-analysis of short-term treatment effects. *Semin Orthod.* 2014;20:324–338.
- Cozza P, Baccetti T, Franchi L, De Toffol L, McNamara JA Jr. Mandibular changes produced by functional appliances in Class II malocclusion: a systematic review. *Am J Orthod Dentofacial Orthop*. 2006;129:599.e1–e12; discussion e1– e6.
- Vogt W. The Forsus Fatigue Resistant Device. J Clinical Orthod. 2006;40:368–377.
- Linjawi AI, Abbassy MA. Dentoskeletal effects of the Forsus[™] fatigue resistance device in the treatment of class II malocclusion: a systematic review and meta-analysis. J Orthod Sc.i 2018;7:5.
- Aslan BI, Kucukkaraca E, Turkoz C, Dincer M. Treatment effects of the Forsus Fatigue Resistant Device used with miniscrew anchorage. *Angle Orthod*. 2014;84:76–87.
- Elkordy SA, Abouelezz AM, Fayed MM, et al. Threedimensional effects of the mini-implant-anchored Forsus Fatigue Resistant Device: a randomized controlled trial. *Angle Orthod*. 2016;86:292–305.
- Elkordy SA, Abouelezz AM, Fayed MMS, Aboulfotouh MH, Mostafa YA. Evaluation of the miniplate-anchored Forsus Fatigue Resistant Device in skeletal Class II growing subjects: a randomized controlled trial. *Angle Orthod.* 2019;89:391–403.
- Gandedkar NH, Shrikantaiah S, Patil AK, et al. Influence of conventional and skeletal anchorage system supported fixed functional appliance on maxillo-mandibular complex and temporomandibular joint: a preliminary comparative cone beam computed tomography study. *Int Orthod.* 2019;17: 256–268.
- Turkkahraman H, Eliacik SK, Findik Y. Effects of miniplate anchored and conventional Forsus Fatigue Resistant Devices in the treatment of Class II malocclusion. *Angle Orthod.* 2016;86:1026–1132.

- Eissa O, Ei-Shennawy M, Gaballah S, El-Meehy G, El Bialy T. Treatment outcomes of Class II malocclusion cases treated with miniscrew-anchored Forsus Fatigue Resistant Device: a randomized controlled trial. *Angle Orthod.* 2017; 87:824–833.
- Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019; 366:14898.
- 15. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919.
- Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol.* 2011;64:383–394.
- Guyatt GH, Oxman AD, Vist G, et al. GRADE guidelines: 4. Rating the quality of evidence—study limitations (risk of bias). *J Clin Epidemiol.* 2011;64:407–415.
- Guyatt GH, Oxman AD, Montori V, et al. GRADE guidelines:
 Rating the quality of evidence—publication bias. *J Clin Epidemiol.* 2011;64:1277–1282.
- Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines 6. Rating the quality of evidence—imprecision. *J Clin Epidemiol.* 2011;64:1283–1293.
- 20. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 7. Rating the quality of evidence—inconsistency. *J Clin Epidemiol.* 2011;64:1294–1302.
- Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 8. Rating the quality of evidence—indirectness. *J Clin Epidemiol.* 2011;64:1303–1310.
- 22. Hajeer MY, Millett DT, Ayoub AF, Siebert JP. Applications of 3D imaging in orthodontics: part I. *J Orthod.* 2004;31:62–70.
- Al-Abdwani R, Moles DR, Noar JH. Change of incisor inclination effects on points A and B. *Angle Orthod.* 2009;79: 462–467.
- 24. Linjawi AI, Abbassy MA. Dentoskeletal effects of the Forsus Fatigue Resistance Device in the treatment of class II malocclusion: a systematic review and meta-analysis. *J Orthod Sci.* 2018;7:5.